arXiv:0811.3169v1 [math.AG] 19 Nov 2008

Tempered fundamental group and metric graph of a Mumford curve

Emmanuel Lepage

November 19, 2008

Contents

1 Tempered fundamental group 4
- 1.1 Definition 4
- 1.2 Mochizuki's results on the pro-(p') tempered group of a curve . . 7

2 Abelianized tempered fundamental group of a curve 9
- 2.1 Tempered fundamental group of an abelian variety 9
- 2.2 Jacobian variety and $\pi_1^{\mathrm{temp}^{\mathrm{ab}}}$ of a curve 10

3 Alterations and tempered fundamental group 12
- 3.1 Preliminaries about the skeleton of a Berkovich space 12
- 3.2 Invariance of π_1^{temp} by base change of algebraically closed fields . 15
- 3.3 Products and tempered fundamental group 16

4 Metric structure of the graph of the stable model of a curve 17
- 4.1 Preliminaries 19
  - 4.1.1 Cusps 20
  - 4.1.2 Vertices of the skeleton 20
- 4.2 Case of $\mathbf{P}^1\backslash\{z_1,\dots,z_n\}$ 21
- 4.3 Case of a punctured elliptic curve 22
- 4.4 Case of a Mumford curve 25
  - 4.4.1 Reminder on Mumford curves and currents 25
  - 4.4.2 Preliminary results on the ramifications of torsors corresponding to currents 27
  - 4.4.3 The metric graph of the reduction and the tempered fundamental group 28

A A combinatorial result 31

Introduction

This paper is an attempt to give some general results on the tempered fundamental group of p-adic smooth algebraic varieties.
The tempered fundamental group was introduced in [2, part III] as a sort of analog of the topological fundamental group of complex algebraic varieties; its

profinite completion coincides with Grothendieck's algebraic fundamental group, but it has itself many infinite discrete quotients in general.
Since the analytification (in the sense of Berkovich or of rigid geometry) of a finite étale covering of a p-adic varieties is not necessarily a topological covering, André had to use a slightly larger notion of covering. He defines tempered coverings, which are étale coverings in the sense of de Jong (that is to say that locally on the Berkovich topology, it is a direct sum of finite coverings) such that, after pulling back by some finite étale covering, they become topological coverings (for the Berkovich topology). Then the tempered fundamental group is a prodiscrete group that classifies those tempered coverings. To give a more handful description, if one has a sequence of pointed finite coverings $((S_i,s_i))_{i\in\mathbf{N}}$ such that the corresponding pointed pro-covering of (X,x) is the universal pro-covering of (X,x), and if (S_i^∞,s_i^∞) is a universal topological covering of S_i, the tempered fundamental group of X can be seen as $\pi_1^{\text{temp}}(X,x)=\varprojlim_i \text{Gal}(S_i^\infty/X)$. Therefore, to understand the tempered fundamental group of a variety, one mainly has to understand the topological behavior of the finite étale coverings of this variety.
There are many differences between the tempered fundamental group in the p-adic case and the topological fundamental group in the complex case. First, the tempered fundamental group is not discrete in general. It is also much more difficult to describe explicitly (such a description is available for elliptic curves, or more generally abelian varieties as will be studied in this article). As proved in [13] (and recalled in part 1.2), the tempered fundamental group of a curve depends hugely on the combinatorial structure of its stable reduction (this suggests a geometric anabelian behavior of the tempered fundamental group which as no algebraic or complex counterpart). On the other hand, the tempered fundamental group, like the algebraic fundamental group, is also defined over non algebraically closed nonarchimedean field, and thus it interacts interestingly with Galois theory (which gives it a number theoretical interest). However, in this article, we will only study the geometric tempered fundamental group. We will prove here some results for the tempered fundamental group which are classical results for the profinite fundamental group or the topological fundamental group of complex varieties: we will link the abelianized tempered fundamental group of a curve to the tempered fundamental group of its Jacobian variety and we will show a Künneth formula for the product of manifolds. But, to illustrate that it depends much more on the variety itself than the profinite or the complex fundamental group, we will show that one can recover the metric structure of the graph of the stable model of a Mumford curve from the tempered fundamental group.

After recalling the basic results on the tempered fundamental group given in [2] (and deduce from them the birational invariance of the tempered fundamental group in proposition (1.6)), we will recall the results of Mochizuki concerning the tempered fundamental group of a curve and the stable reduction of this curve: whereas the profinite fundamental group or the topological fundamental group of a smooth curve of type (g,n) only depends in the complex case on g and n, the tempered fundamental group of a curve depends much more on the curve itself. How much is an unsolved problem, which is the leading thread of the present paper.
Mochizuki thus proved in [13] that one could recover from the tempered fun-

damental group of a curve the graph of its stable reduction (theorem (1.7)), even in the pro-(p') case (we will denote by $\pi_1^{\text{temp}}(X)^{(p')}$ the group classifying coverings that became topological after base change by a finite Galois covering of order prime to p).
More precisely, the vertices correspond to the classes of maximal subgroups of $\pi_1^{\text{temp}}(X)^{(p')}$ (thus called verticial subgroup) and the edges to non trivial classes of intersection of verticial subgroups.

We will then study the tempered fundamental group of an abelian variety in the geometric case. We will be able to give an explicit description of the tempered fundamental group, as was already done for elliptic curves in [2, III.2.3.2]. The point is that the finite étale coverings are well understood (they are abelian varieties, and they are dominated by the multiplication by n on A for some n), and the topological coverings are well understood too, thanks to the p-adic uniformization of abelian varieties.
We will deduce from this that $\pi_1^{\text{temp}}(A)$ is isomorphic to $\widehat{\mathbf{Z}}^{2g-d} \times \mathbf{Z}^d$ (see part 2.1), where g is the dimension of A and d is the rank of the topological fundamental group of A (which is also the dimension of the toric part of the semistable reduction).
One can then prove that, just as in the profinite case or the topological case, the abelianized tempered fundamental group of a curve X is canonically isomorphic to the tempered fundamental group of its Jacobian variety A (theorem (2.1)). The proof combines the birational morphism $X^g/\mathfrak{S}_g \to A$, the birational invariance of the tempered fundamental group, and the Künneth formula that will be proved in the next part.

For more general cases, one does not have such nice descriptions of the topological behavior of a smooth variety, not to mention the topological behavior of all its finite etale coverings. Nevertheless, thanks to de Jong's alteration results (see [6] and [7]) and to Berkovich's work (see [3], recalled in part 3.1) on the topological structure of the generic fiber of a polystable formal scheme over the integral ring of a complete nonarchimedean field, one can prove that some Zariski open dense subset U of the smooth scheme X is homotopically equivalent to a certain polysimplicial set given by the combinatorial structure of the special fiber of some alteration of the variety.
One can then deduce from the results of [3] that for any isommetric embedding of K in an algebraically closed field K', $|U_{K'}| \to |U|$ is a homotopy equivalence. Knowing this, and using a proposition of André which says that $U \to X$ induces an isomorphism on topological groups, we will be able to prove the invariance of the tempered fundamental group of a smooth scheme under algebraically closed extensions (proposition (3.7).
In the same way, if V is a Zariski dense open subset of another smooth scheme Y satisfying the same properties as U, one can deduce from the results of [3] that $|U \times V| \to |U| \times |V|$ is a homotopy equivalence. Thanks to this, we will prove a Künneth formula for the tempered fundamental group of smooth schemes (proposition (3.8)).

Finally, we will prove in theorem (4.10) that, in the case of a Mumford curve, one may in fact recover the length of the edges from $\pi_1^{\text{temp}}(X)$ (obviously, one needs the whole $\pi_1^{\text{temp}}(X)$ and not only the pro-(p') version which only depends of the pro-(p') graph of groups of the stable reduction; thus this can only be done when $p \neq 0$, contrarily to Mochizuki's result).

We will deduce from Mochizuki's results that one can decide from the tempered fundamental group of a curve whether a finite covering is split over some vertex of the curve (and in a similar way, one can decide whether it is ramified over some cusp).

Knowing this, we will start by proving this for a punctured projective line and a punctured elliptic curve, to give an insight of the method in simpler cases, where the coverings we will use are described explicitly: for a punctured projective line, we will concentrate on coverings like $(\ )^{p^e} : \mathbf{G}_m \overset{z \mapsto z^{p^e}}{\to} \mathbf{G}_m$ (one can in this case easily calculate explicitly the number of preimage of a point of the Berkovich projective line), and for elliptic curves on coverings obtained by patching of those types of coverings.

In the more general case of a Mumford curve $X = \Omega/\Gamma$, we will study coverings of the topological uniformization Ω that descend to some finite topological covering of X that behave like $(\ )^{p^e}$ over big affinoids of Ω. More precisely, every covering of Ω which descends to X (in some non unique way) is the pullback of $(\ )^{p^e}$ along a Γ-equivariant invertible section of Ω, and conversely. The equivariance prevents us to simply consider a homography, but we can consider a section of Ω which is arbitrarily close, on a given affinoid subset of Ω, to an homography. This will be enough to ensure that this covering is split over the same vertices of a big affinoid subset of Ω as the one obtained by pulling back $(\ )^{p^e}$ along the homography.

Using such constructions, one can recover from the tempered fundamental group the length of any loop of any topological covering of X. In order to get the whole metric structure of the graph of the stable model of X, we will end this article by proving a purely combinatorial result that shows that, if one knows the length of every loop of every covering of a graph whose edges have valency ≥ 3, one knows the length of every edge (proposition (A.1)).

The recovering of the metric graph of the stable model from the tempered fundamental group can be quite easily extended to the case of punctured Mumford cases, but the proof does not seem to extend easily to more general curves, as one cannot find that easily coverings of order divisible by p where one can say much about the graph of the stable reduction.

The proofs here also only consider some very simple coverings (for which we do not even describe completely the graph of the stable model), and it is hard to imagine what can recover from the whole tempered fundamental group.

This work is part of a PhD thesis. I would like to thank my advisor, Yves André, for guiding my work, suggesting the main idea of some proofs presented here, and taking the time of reading and correcting this work.

1 Tempered fundamental group

1.1 Definition

Let K be a complete nonarchimedean field.

Following [1, §4], a *K-manifold* will be a smooth paracompact strictly K-analytic space. For example, if X is a smooth algebraic K-variety, X^{an} is a

K-manifold (and in fact, we will mainly be interested in those spaces). Then, thanks to [3], such a manifold is locally contractible (we will explain in more detail the results of [3] in section 3.1).

Recall from [5] that a morphism $f : S' \to S$ is said to be an *étale covering* if S is covered by open subsets U such that $f^{-1}(U) = \coprod V_j$ and $V_j \to U$ is étale finite.

Then, André defines tempered coverings as follows:

Definition 1.1 *(*[2, def. 2.1.1]*) An étale covering $S' \to S$ is* tempered *if it is a quotient of the composition of a topological covering $T' \to T$ and of a finite étale covering $T \to S$.*

This is equivalent to say that it becomes a topological covering after pullback by some finite étale covering.

We denote by $\mathrm{Cov}^{\mathrm{temp}}(X)$ the category of tempered coverings of X (with the obvious morphisms).

Let $\bar{x}$ be a geometric point of X. Then one has a functor

$$F_{\bar{x}} : \mathrm{Cov}^{\mathrm{temp}}(X) \to \mathrm{Set}$$

which maps a covering $S \to X$ to the set $S_{\bar{x}}$. If $\bar{x}$ and $\bar{x}'$ are two geometric points, then, according to [5, prop 2.9], $F_{\bar{x}}$ and $F_{\bar{x}'}$ are (non canonically) isomorphic.

The tempered fundamental group of X pointed at $\bar{x}$ is

$$\pi_1^{\mathrm{temp}}(X, \bar{x}) = \mathrm{Aut}\, F_{\bar{x}}.$$

This is a prodiscrete topological group, for which the stabilizers $(\mathrm{Stab}_{F(S)}(s))_{S \in \mathrm{Cov}^{\mathrm{temp}}(X), s \in F_{\bar{x}}(S)}$ form a basis of open subgroups of $\pi_1^{\mathrm{temp}}(X, \bar{x})$. The tempered fundamental group depends on the basepoint only up to inner automorphism (this topological group, considered up to conjugation, will sometimes be denoted simply $\pi_1^{\mathrm{temp}}(X)$).

The full subcategory of tempered coverings S for which $F_{\bar{x}}(S)$ is finite is equivalent to $\mathrm{Cov}^{\mathrm{alg}}(S)$, so that

$$\widehat{\pi_1^{\mathrm{temp}}(X, \bar{x})} = \pi_1^{\mathrm{alg}}(X, \bar{x})$$

(where $\widehat{\ }$ denotes here, and in the sequel, the profinite completion).

For any morphism $X \to Y$, the pullback defines a functor $\mathrm{Cov}^{\mathrm{temp}}(Y) \to \mathrm{Cov}^{\mathrm{temp}}(X)$. If $\bar{x}$ is a geometric point of X with image $\bar{y}$ in Y, this gives rise to a continuous homomorphism

$$\pi_1^{\mathrm{temp}}(X, \bar{x}) \to \pi_1^{\mathrm{temp}}(Y, \bar{y})$$

(hence an outer morphism $\pi_1^{\mathrm{temp}}(X) \to \pi_1^{\mathrm{temp}}(Y)$).

One has the analog of the usual Galois correspondence:

Theorem 1.2 ([2, Th. 1.4.5]) *$F_{\bar{x}}$ induces an equivalence of categories between the category of direct sums of tempered coverings of X and the category $\pi_1^{\mathrm{temp}}(X, \bar{x}) - \mathrm{Set}$ of discrete sets endowed with a continuous left action of $\pi_1^{\mathrm{temp}}(X, \bar{x})$.*

If S is a finite Galois covering of X, its universal topological covering S^∞ is still Galois and every connected tempered covering is dominated by such a Galois tempered covering.
If $((S_i,\bar{s}_i))_{i\in\mathbf{N}}$ is a cofinal projective system (with morphisms $f_{ij}:S_i\to S_j$ which maps s_i to s_j for $i\geq j$) of geometrically pointed Galois finite étale coverings of $(X,\bar{x})$, let $((S_i^\infty,\bar{s}_i^\infty))_{i\in\mathbf{N}}$ be the projective system, with morphisms f_{ij}^∞ for $i\geq j$, of its pointed universal topological coverings. Then $F_{\bar{x}}(S_i^\infty)=\pi_1^{\mathrm{temp}}(X,\bar{x})/\operatorname{Stab}_{F(S_i^\infty)}(\bar{s}_i^\infty)$ is naturally a quotient group of $\pi_1^{\mathrm{temp}}(X,\bar{x})$ for which s_i^∞ is the neutral element. Moreover G acts by G-automorphisms on $F_{\bar{x}}(S_i^\infty)$ by right translation (and thus on S_i^∞ thanks to the Galois correspondence of theorem (1.2)). Thus one gets a morphism $\pi_1^{\mathrm{temp}}(X,\bar{x})\to\operatorname{Gal}(S_i^\infty/X)$.
As $f_{ij}^\infty(s_i^\infty)=s_j^\infty$, these morphisms are compatible with $\operatorname{Gal}(S_i^\infty/X)\to\operatorname{Gal}(S_j^\infty/X)$.
Then, thanks to [2, lemma III.2.1.5],

Proposition 1.3

$$\pi_1^{\mathrm{temp}}(X,\bar{x})\to\varprojlim\operatorname{Gal}(S_i^\infty/X)$$

is an isomorphism.

We will also have to use the following result by André:

Proposition 1.4 ([2, prop. III.1.1.4]) *Let $\overline{S}$ be a manifold, and let Z be a Zariski closed nowhere dense reduced analytic subset. Then any topological covering of $S:=S\backslash Z$ extends uniquely to a topological covering. Thus $\pi_1^{\mathrm{top}}(S,s)\to\pi_1^{\mathrm{top}}(\overline{S},s)$ is an isomorphism.*

We can deduce from this that:

Proposition 1.5 ([2, th. III.2.1.11, prop. III.2.1.13]) *Assume K algebraically closed. Let $\overline{S}$ be a manifold, and let Z be a Zariski closed nowhere dense reduced analytic subset. Then the functor from tempered coverings of $\overline{S}$ to tempered coverings of $S=\overline{S}\backslash Z$ is fully faithful. If Z is of codimension ≥ 2, this functor is an equivalence of categories.*

We will follow the proof of [10, cor. X.3.4] to get the birational invariance of the tempered fundamental group of smooth and proper K-schemes.
Recall that every rational map between smooth and proper K-schemes is defined above the complement of a closed subset of codimension ≥ 2.
Let $f:X\to Y$ is a dominant rational map between smooth and proper K-schemes. f is defined on a Zariski open subset U of X (denote by f_U the morphism $U\to Y$ and by i_U the immersion $U\to X$) whose complement is of codimension ≥ 2, one gets a functor from $\mathrm{Cov}^{\mathrm{temp}}(Y)$ to $\mathrm{Cov}^{\mathrm{temp}}(X)$ and one can compose it with a quasi-inverse of $\mathrm{Cov}^{\mathrm{temp}}(U)\to\mathrm{Cov}^{\mathrm{temp}}(X)$: one thus get a functor $f_{(U)}^*:\mathrm{Cov}^{\mathrm{temp}}(Y)\to\mathrm{Cov}^{\mathrm{temp}}(X)$, such that $i_U^*f_{(U)}^*$ is isomorphic to f_U^*. If one takes another Zariski open subset U' of X satisfying the same properties, one gets that $i_{U\cap U'}^*f_{(U)}^*$ and $i_{U\cap U'}^*f_{(U')}^*$ are both isomorphic to $f_{U\cap U'}^*$, and thus $f_{(U)}^*$ and $f_{(U')}^*$ are isomorphic, since $X\backslash U\cap U'$ is also of codimension ≥ 2. Thus one gets an outer homomorphism of topological groups $f_*:\pi_1^{\mathrm{temp}}(X)\to\pi_1^{\mathrm{temp}}(Y)$, which does not depend on U. In particular if f is a morphism of schemes, one can choose $U=X$ and thus f_* is the usual outer morphism $\pi_1^{\mathrm{temp}}(X)\to\pi_1^{\mathrm{temp}}(Y)$.

Let $g : Y \to Z$ be another dominant rational map between smooth and proper K-scheme, it is defined on a Zariski open subset V of Y of codimension ≥ 2, and $gf : X \to Z$ is also a dominant rational map between smooth and proper schemes, so it is also defined over a Zariski open subset W of X of codimension ≥ 2. Let $U_0 = U \cap f_U^{-1}(V) \cap W$ ($X \backslash U_0$ does not need to be of codimension ≥ 2). There are morphisms $U_0 \to V$ and $V \to Z$ representing f and g such that the composed morphism $(gf)_{U_0} : U_0 \to Z$ represents gf. One then gets that $i^*_{U_0} f^*_{(U)} g^*_{(V)}$ and $i^*_{U_0} (gf)^*_{(W)}$ are both isomorphic to $(gf)^*_{U_0}$. Since $i^*_{U_0}$ is fully faithful, $f^*_{(U)} g^*_{(V)}$ and $(gf)^*_{(W)}$ are isomorphic (and $g_* f_* = (gf)_*$). Thus one gets a functor from the category of smooth and proper K-schemes with dominant rational maps to the category of groups with outer homomorphisms. In particular,

Proposition 1.6 *Let $X \to Y$ be a birational map between smooth and proper K-schemes. Then*

$$\pi_1^{\mathrm{temp}}(X) \to \pi_1^{\mathrm{temp}}(Y)$$

is an isomorphism.

If one considers the full subcategory $\mathrm{Cov}^{\mathrm{temp}}(X)^{(p')}$ of tempered coverings that become topological after pullback by a finite Galois covering of order prime to p where p is the residual characteristic, one gets in the same fashion a pro-(p') version $\pi_1^{\mathrm{temp}}(X, \bar{x})$ of the tempered fundamental group (see [13, Rm. 3.10.1.] in the case of a curve).

1.2 Mochizuki's results on the pro-(p') tempered group of a curve

Following [12, Appendix], a *semigraph* $\mathbb{G}$ will be given by a set $\mathcal{V}$ of vertices, a set $\mathcal{E}$ of edges and, for every $e \in \mathcal{E}$ a set with cardinality ≤ 2 of branches $\mathcal{B}_e$ with a $\zeta_e : \mathcal{B}_e \to \mathcal{V}$ (we will say that a branch b of e *abuts* to v if $\zeta_e(b) = v$). We will say that a semigraph is a *graph* if for every $e \in \mathcal{E}$, e has exactly two branches.

Recall from [13, def. 2.1] that if one has a semigraph $\mathcal{G}$, a structure of *semigraph of anabelioid* on this graph corresponds to the following data: for each vertex or edge x, a Galois category (also named connected anabelioid in [13]) $\mathcal{G}_x$, and for each branch b of edge e abutting to a vertex v, a morphism of anabelioids (*i.e.* an exact functor) $b_* : \mathcal{G}_e \to \mathcal{G}_v$). Recall also that working with Galois categories is equivalent to working with profinite groups up to inner automorphism.

A *covering* S of $\mathcal{G}$ consists of data (S_v, ϕ_e), where for a vertex v, S_v is an object of $\mathcal{G}_v^{\top}$ (the category of arbitrary disjoint unions of connected objects of $\mathcal{G}_v$, that is the topos $\pi_1(\mathcal{G}_v)-\mathrm{Set}$), and for every edge e with branches b_1 and b_2 abutting to v_1 and v_2 ϕ_e is an isomorphism between $b_{1*}S_{v_1}$ and $b_{2*}S_{v_2}$. One has a natural notion of morphism of such coverings, so that one gets a category $\mathcal{B}^{\mathrm{cov}}(\mathcal{G})$. Mochizuki associates canonically to such a covering a semigraph of anabelioids $\mathcal{G}'$ above $\mathcal{G}$.

An object of $\mathcal{B}^{\mathrm{cov}}(\mathcal{G})$ is finite if each S_v is in $\mathcal{G}_v$, *topological* if for each v S_v is a constant object of $\mathcal{G}_v^{\top}$, and *tempered* if it becomes topological after pulling back along some finite covering. The full subcategory of tempered coverings is then denoted by $\mathcal{B}^{\mathrm{temp}}(\mathcal{G})$ (and $\mathcal{B}^{\mathrm{alg}}(\mathcal{G})$ denotes the full subcategory of finite

coverings; if $\mathbb{G}$ is connected, then it is a Galois category of fundamental group denoted as $\pi_1^{\mathrm{alg}}(\mathcal{G})$).
Then, if v is a vertex of $\mathbb{G}$ and F is a fundamental functor of $\mathcal{G}_v$ (and thus extends to a point of $\mathcal{G}_v^{\top}$), one can define a functor $F_{(v,F)} : \mathcal{B}^{\mathrm{cov}}(\mathcal{G}) \to \mathrm{Set}$ which maps S to $F(S_v)$ (if one changes the base point (v, F), one gets an isomorphic functor), and let $F_{(v,F)}^{\mathrm{temp}}$ be its restriction to $\mathcal{B}^{\mathrm{temp}}(\mathcal{G})$. Then we define $\pi_1^{\mathrm{temp}}(\mathcal{G}, (v, F))$ to be $\mathrm{Aut}(F_{(v,F)}^{\mathrm{temp}})$.

Let $(\overline{X}, D)$ be a smooth n-pointed curve of type g over a complete discrete valuation field K, let $X = \overline{X} \backslash D$, let $(\overline{\mathcal{X}}, \mathcal{D})$ be a semistable model over $O_{\overline{K}}$, where $\overline{K}$ is the completion of an algebraic closure of K, and let $\mathcal{X} = \overline{\mathcal{X}} \backslash \mathcal{D}$.
The semigraph $\mathbb{G}_{\mathcal{X}}^{c}$ of $\mathcal{X}_s$ is defined as follows: the vertices are the irreducible components of $\mathcal{X}_s$, the edges are the nodes and the marked points. A node e has two branches that abut to the irreducible components that contain e; a marked point e has only one branch that abuts to the irreducible component containing the marked point.
When $\mathcal{X}$ is the stable model of X, $\mathbb{G}_{\mathcal{X}}^{c}$ will simply be denoted by $\mathbb{G}_X^c$ (or $\mathbb{G}^c$ when there is no risk of confusion).
One can endow the semigraph $\mathbb{G}^c$ of the special fiber $\mathcal{X}_s$ of $\mathcal{X}$ with a structure of "semigraph of anabelioids" $\mathcal{G}^c$. Indeed for a vertex v_i corresponding to an irreducible component C_i of $\mathcal{X}_s$, consider the open subset U_i of the normalization $\overline{C}_i'$ of $\overline{C}_i$ which is the complementary of the marked points and of the preimages of the double points of $\mathcal{X}_s$ (the points of $\overline{C}_i' - U_i$ thus correspond exactly to branches abutting to v_i). Then the group $\Pi_{(v_i)}$ is the tame fundamental group $\pi_1^{\mathrm{t}}(U_i)$ of U_i in $\overline{C}_i'$. The group of an edge is $\widehat{\mathbf{Z}(1)}^{(p')} = \varprojlim_{(n,p)=1} \mu_n (\simeq \widehat{\mathbf{Z}}^{(p')})$ ((p') is for the pro-prime-to-p maximal quotient), which is canonically isomorphic to the subgroup of monodromy in $\pi_1^{\mathrm{t}}(U_i)$ of a point in $\overline{C}_i' - U_i$. The morphism corresponding to a branch is the embedding of the monodromy group of the corresponding point of $\overline{C}_i' - U_i$ (defined up to conjugation), whereas, for an edge with two branches, one identifies the two $\widehat{\mathbf{Z}(1)}^{(p')}$ by $x \mapsto x^{-1}$.
If $\mathcal{G}^{(p')}$ denotes the semigraph of anabelioids obtained from $\mathcal{G}$ by replacing each profinite group by its pro-(p') completion, and if $\pi_1^{\mathrm{temp}}(X_{\overline{K}})^{(p')}$ is the pro-(p') version of the tempered fundamental group of $X_{\overline{K}}$,

$$\pi_1^{\mathrm{temp}}(\mathcal{G}_X^{c,(p')}) = \pi_1^{\mathrm{temp}}(X_{\overline{K}})^{(p')}.$$

Mochizuki then shows:

Theorem 1.7 ([13, cor. 3.11]) *If X_α and X_β are two curves, every isomorphism $\gamma : \pi_1^{\mathrm{temp}}(X_{\alpha,\overline{K}}) \simeq \pi_1^{\mathrm{temp}}(X_{\beta,\overline{K}})$ determines, functorially in γ up to 2-isomorphism, an isomorphism of semigraphs of anabelioids $\gamma' : \mathcal{G}_{X_\alpha}^{c} \simeq \mathcal{G}_{X_\beta}^{c}$.*

More precisely, the following induced diagram of topological group is commutative:

$$\begin{tikzcd}
\pi_1^{\mathrm{temp}}(\mathcal{G}_{X_\alpha}^{c,(p')}) \arrow[r, "\pi_1^{\mathrm{temp}}(\gamma'^{(p')})"] \arrow[d, equal] & \pi_1^{\mathrm{temp}}(\mathcal{G}_{X_\beta}^{c,(p')}) \arrow[d, equal] \\
\pi_1^{\mathrm{temp}}(X_{\alpha,\overline{K}})^{(p')} \arrow[r, "\gamma^{(p')}"] & \pi_1^{\mathrm{temp}}(X_{\beta,\overline{K}})^{(p')}
\end{tikzcd}$$

The vertices of the graph then correspond to the conjugacy classes of maximal compact subgroups of $\pi_1^{\mathrm{temp}}(X)^{(p')}$ (thus called *verticial* subgroups of $\pi_1^{\mathrm{temp}}(X)^{(p')}$), and edges to conjugacy class of nontrivial intersections of verticial subgroups.

2 Abelianized tempered fundamental group of a curve

Here, we will be interested in the tempered fundamental group of an abelian variety and in the abelianized tempered fundamental group of a curve.
We will first prove that if A is an abelian variety over an algebraically closed complete nonarchimedean field of characteristic 0, the tempered fundamental group is abelian and fits in a split exact sequence:

$$0 \to T \to \pi_1^{\mathrm{temp}}(A) \to \pi_1^{\mathrm{top}}(A) \to 0$$

where T is profinite (and more precisely isomorphic to $\widehat{\mathbf{Z}}^n$ for some n).
Then we will prove that (as in the case of algebraic fundamental groups or complex topological fundamental groups) if C is a curve and A its Jacobian variety, the natural morphism $\pi_1^{\mathrm{temp}}(C)^{\mathrm{ab}} \to \pi_1^{\mathrm{temp}}(A)$ is an isomorphism.

2.1 Tempered fundamental group of an abelian variety

Let K be an algebraically closed complete nonarchimedean field of characteristic 0.
Let A be an abelian variety over K, and let g be its dimension.

Recall the basics of p-adic uniformization of abelian varieties.
By [9], there is a commutative algebraic group G (more precisely a semiabelian variety) and a surjective analytic morphism $u : G^{\mathrm{an}} \to A^{\mathrm{an}}$ which is the universal topological covering of A and $\ker u$ is a discrete free abelian group Λ of rank d. Moreover, we know that $(A^{(n)} \to A)_{n\in\mathbf{N}}$ ($\mathbf{N}$ being ordered by divisibility), with $A^{(n)}$ a copy of A and $A^{(n)} \to A$ multiplication by n, is a cofinal family of finite Galois coverings of A (by [10, lecture XI]).
Let $G^{(n)}$ be the universal topological covering of $A^{(n)}$ (which is isomorphic to G since $A^{(n)}$ is isomorphic to A): one has

$$\pi_1^{\mathrm{temp}}(A) = \varprojlim_n \mathrm{Gal}(G^{(n)}/A).$$

Since $\pi_1^{\mathrm{top}}(A^{(n)}) = \pi_1^{\mathrm{top}}(A) = \Lambda \simeq \mathbf{Z}^d$ is residually finite and is a subgroup of finite index of $\mathrm{Gal}(G^{(n)}/A)$, $\mathrm{Gal}(G^{(n)}/A)$ is residually finite for every n, so $\pi_1^{\mathrm{temp}}(A)$ is also topologically residually finite as a projective limit of residually finite groups. Thus $\pi_1^{\mathrm{temp}}(A) \to \pi_1^{\mathrm{alg}}(A)$ is injective and, since $\pi_1^{\mathrm{alg}}(A)$ is abelian, $\pi_1^{\mathrm{temp}}(A)$ is also an abelian group.

If $n|m$, one has the following commutative diagram :

$$\begin{tikzcd}
0 \arrow[r] & \mathrm{Gal}(G^{(m)}/G) \arrow[r] \arrow[d] & \mathrm{Gal}(G^{(m)}/A) \arrow[r] \arrow[d] & \mathrm{Gal}(G/A) = \Lambda \arrow[r] \arrow[d, equal] & 0 . \\
0 \arrow[r] & \mathrm{Gal}(G^{(n)}/G) \arrow[r] & \mathrm{Gal}(G^{(n)}/A) \arrow[r] & \mathrm{Gal}(G/A) = \Lambda \arrow[r] & 0
\end{tikzcd}$$

Let us write $T(G) = \varprojlim \mathrm{Gal}(G^{(n)}/G)$. This is a profinite abelian group, so it splits (canonically) as the product of its pro-p-Sylow subgroups $T_l(G)$.
By taking the projective limit in the previous commutative diagram, one gets the following axact sequence (it is right exact because the $\mathrm{Gal}(G^{(n)}/G$ are finite):

$$\begin{tikzcd}
0 \arrow[r] & T(G) \arrow[r] & \pi_1^{\mathrm{temp}}(A) \arrow[r] & \Lambda \arrow[r] & 0 ,
\end{tikzcd}$$

with $\pi_1^{\mathrm{temp}}(A)$ abelian. Thus it is an exact sequence of abelian groups.
But Λ is a free abelian group, so the exact sequence must be split. One thus gets a non canonical isomorphism:

$$\pi_1^{\mathrm{temp}}(A) \simeq \Lambda \times T(G).$$

By taking the pro-l completion of the isomorphism here above, one gets:

$$\mathbf{Z}_l^{2g} \simeq \pi_1^{\mathrm{alg}}(A)^l \simeq \widehat{\pi_1^{\mathrm{temp}}(A)}^{l} \simeq \mathbf{Z}_l^d \times T_l(G),$$

and so $T_l(G) \simeq \mathbf{Z}_l^{2g-d}$. We finally obtained a non canonical isomorphism:

$$\pi_1^{\mathrm{temp}}(A) \simeq \mathbf{Z}^d \times \widehat{\mathbf{Z}}^{2g-d}.$$

2.2 Jacobian variety and $\pi_1^{\mathrm{temp}^{\mathrm{ab}}}$ of a curve

If G is a topological prodiscrete group with a countable basis of neighborhoods of 1, G^{ab} will be the topological group $G/\overline{D(G)}$ (where $\overline{D}(G)$ is the closure of the derived subgroup of G), it is also a prodiscrete group and $G \to G^{\mathrm{ab}}$ makes $G^{\mathrm{ab}} - \mathrm{Ens}$ as a full subcategory of $G - \mathrm{Ens}$.

Let K be a complete discrete valuation field of characteristic 0, let $\bar{K}$ be the completion of its algebraic closure, let C be a curve over K and let A be the Jacobian variety of $C_{\bar{K}}$.
Let P be a closed point of $C_{\bar{K}}$.
Consider the morphism $C_{\bar{K}} \to A$ that maps x to the divisor $[x] - [P]$. One gets a homomorphism $\pi_1^{\mathrm{temp}}(C_{\bar{K}}, P)^{\mathrm{ab}} \to \pi_1^{\mathrm{temp}}(A, 0)$ that factorizes through $\pi_1^{\mathrm{temp}}(C_{\bar{K}}, P)^{\mathrm{ab}}$ since $\pi_1^{\mathrm{temp}}(A, 0)$ is abelian.

Theorem 2.1 *The morphism $\pi_1^{\mathrm{temp}}(C_{\bar{K}}, P)^{\mathrm{ab}} \to \pi_1^{\mathrm{temp}}(A, 0)$ is an isomorphism.*

Proof : We have a morphism $C_{\bar{K}}^g \to A$ which maps $(x_1, \dots, x_g)$ to the divisor $[x_1] + \cdots + [x_g] - g[P]$ of $C_{\bar{K}}$. This morphism is invariant by the action of $\mathfrak{S}_g$ on

$C^g_{\bar K}$ and thus factorizes in a morphism $C^{(g)}_{\bar K} := C^g_{\bar K}/\mathfrak{S}_g \to A$. Remind that this is a birational morphism and that $C^{(g)}_{\bar K}$ is smooth over $\bar K$ (voir [11, Th. 5.1.(a), prop. 3.2]).
We thus get a sequence of morphisms:

$$C_{\bar K} \to C^g_{\bar K} \to C^{(g)}_{\bar K} \to A,$$

where the left morphism maps x to $(x, P, \dots, P)$ and where the composed morphism is the morphism that maps x to $[x] - [P]$.

Since $C^{(g)}_{\bar K} \to A$ is a birational morphism of proper smooth $\bar K$-varieties, $\pi_1^{\mathrm{temp}}(C^{(g)}_{\bar K}, (P, \dots, P)) \to \pi_1^{\mathrm{temp}}(A, 0)$ is an isomorphism according to proposition (1.6).
Thus $\pi_1^{\mathrm{temp}}(C^{(g)}_{\bar K}, (P \dots, P))$ is abelian and residually finite.
Thus $\pi_1^{\mathrm{temp}}(C_{\bar K}, P) \to \pi_1^{\mathrm{temp}}(C^{(g)}_{\bar K}, (P, \dots, P))$ factorizes through $\phi : \pi_1^{\mathrm{temp}}(C_{\bar K}, P)^{\mathrm{ab}} \to \pi_1^{\mathrm{temp}}(C^{(g)}_{\bar K}, (P, \dots, P))$.

As $\pi_1^{\mathrm{temp}}(C_{\bar K}, P)^{\mathrm{ab}}$ is a projective limit of abelian groups of finite type (and thus residually finite groups), it must be residually finite too.
The following commutative diagram:

$$\begin{tikzcd}
\pi_1^{\mathrm{temp}}(C_{\bar K}, P)^{\mathrm{ab}} \arrow[r, "\phi"] \arrow[d] & \pi_1^{\mathrm{temp}}(C^{(g)}_{\bar K}, (P, \dots, P)) \arrow[d] \\
\pi_1^{\mathrm{alg}}(C_{\bar K}, P)^{\mathrm{ab}} \arrow[r, "\simeq"] & \pi_1^{\mathrm{alg}}(C^{(g)}_{\bar K}, (P, \dots, P))
\end{tikzcd},$$

whose vertical arrows are injective, shows that ϕ is injective.

One may also put an orbifold structure on $C^{(g)}_{\bar K}$, as being $C^g_{\bar K}/\mathfrak{S}_g$ (then $\pi_1^{\mathrm{orb}}(C^{(g)}_{\overline{K}})$ denotes the corresponding extension of $\mathfrak{S}_g$ by $\pi_1^{\mathrm{temp}}(C^g_{\overline{K}})$, as in [2, prop. III.4.5.8]).
One then has the following commutative diagram whose line is exact:

$$\begin{tikzcd}
1 \arrow[r] & \pi_1^{\mathrm{temp}}(C^g_{\bar K}, (P, \dots, P)) \arrow[r, "i"] \arrow[ddr, "\alpha"'] & \pi_1^{\mathrm{orb}}(C^{(g)}_{\bar K}, (P, \dots, P)) \arrow[r] \arrow[d, "\pi_1"] & \mathfrak{S}_g \arrow[r] & 1 \\
& & \pi_1^{\mathrm{orb}}(C^{(g)}_{\bar K}, (P, \dots, P))^{\mathrm{ab}} \arrow[d, "\pi_2"] & & \\
& & \pi_1^{\mathrm{temp}}(C^{(g)}_{\bar K}, (P, \dots, P)) & &
\end{tikzcd}$$

The morphisms i, π_1 and π_2 are open, thus $\alpha = \pi_2 \pi_1 i$ is open: $\operatorname{Im}\alpha$ is an open subgroup of $\pi_1^{\mathrm{temp}}(C^{(g)}_{\bar K}, (P, \dots, P))$.
But, if it is a strict subgroup, $\pi_1^{\mathrm{temp}}(C^{(g)}_{\bar K}, (P, \dots, P))/\operatorname{Im}\alpha$ is a nontrivial abelian group of finite type and thus has a nontrivial finite quotient, which corresponds to an open subgroup of finite index of $\pi_1^{\mathrm{temp}}(C^{(g)}_{\bar K}, (P, \dots, P))$ which

contains $\operatorname{Im}\alpha$.
But $\hat{\alpha}$ is surjective (since $\pi_1^{\mathrm{alg}}(C_{\bar{K}}) \to \pi_1^{\mathrm{alg}}(C_{\bar{K}}^{(g)})$, which factorizes through $\hat{\alpha}$ is surjective), and thus one gets a contradiction: α is also surjective.

Let us now consider the diagonal hommomorphism $\delta : \pi_1^{\mathrm{temp}}(C_{\bar{K}}, P) \to \pi_1^{\mathrm{temp}}(C_{\bar{K}}^g, (P, \dots, P))$. By identifying $\pi_1^{\mathrm{temp}}(C_{\bar{K}}^g, (P, \dots, P))$ to $\pi_1^{\mathrm{temp}}(C_{\bar{K}}, (P, \dots, P))^g$ thanks to proposition (3.8) that we will prove in the next section, δ may be identified to $\pi_1^{\mathrm{temp}}(C_{\bar{K}}, P) \to \pi_1^{\mathrm{temp}}(C_{\bar{K}}, P)^g : g \mapsto (g, 1, \dots, 1)$.
Thus $\pi_1^{\mathrm{temp}}(C_{\bar{K}}^g, (P, \dots, P))$ is generated by the images of $\sigma \circ \delta$ when σ describes $\mathfrak{S}_g$, and as α is invariant by σ, $\alpha \circ \delta$ is surjective, so $\pi_1^{\mathrm{temp}}(C_{\bar{K}}, P)^{\mathrm{ab}} \to \pi_1^{\mathrm{temp}}(C_{\bar{K}}^{(g)}, (P, \dots, P))$ is also surjective, thus it is bijective.
If U is an open subgroup of $\pi_1^{\mathrm{temp}}(C_{\bar{K}}, P)$, the group generated by the $\sigma(\delta(U))$ is an open subgroup of $\pi_1^{\mathrm{temp}}(C_{\bar{K}}^g, (P, \dots, P))$, thus, as α is open and $\mathfrak{S}_g$-invariant, $\alpha \circ \delta$ is open, thus $\pi_1^{\mathrm{temp}}(C_{\bar{K}}, P)^{\mathrm{ab}} \to \pi_1^{\mathrm{temp}}(C_{\bar{K}}^{(g)}, (P, \dots, P))$ is also open, so it is an isomorphism. □

3 Alterations and tempered fundamental group

In this section we will describe two applications of the theorems of existence of semistable alterations proved by de Jong to the tempered fundamental group. Indeed, Berkovich already showed in [3, §9] how those alterations help building a skeleton, homeomorphic to the geometric realization of a polysimplicial set, on which a Zariski open set of the variety retracts.
We will here deduce from the results of Berkovich that, in characteristic zero, the tempered fundamental group of a smooth algebraic variety is invariant by base change of algebraically closed complete fields, and that the tempered fundamental group of the product of two smooth varieties (over an algebraically closed base) is canonically isomorphic to the product of the tempered fundamental groups of each variety.

3.1 Preliminaries about the skeleton of a Berkovich space

Let k be a complete nonarchimedean field and let k° be its ring of integers.
Recall the definition of a polystable morphism of formal schemes:

Definition 3.1 ([3, def. 1.2],[4, section 4.1]) *Let $\phi : \mathfrak{X} \to \mathfrak{Y}$ be a locally finitely presented morphism of formal schemes.*

(i) ϕ is said to be strictly polystable *if, for every point $y \in \mathfrak{Y}$, there exists an open affine neighborhood $\mathfrak{X}' = \operatorname{Spf}(A)$ of $\phi(y)$ and an open neighborhood $\mathfrak{Y}' \subset \phi^{-1}(\mathfrak{X}')$ of x such that the induced morphism $\mathfrak{Y}' \to \mathfrak{X}'$ goes through an étale morphism $\mathfrak{Y}' \to \operatorname{Spf}(B_0) \times_{\mathfrak{X}'} \cdots \times_{\mathfrak{X}'} \operatorname{Spf}(B_p)$ where each B_i is of the form $A\{T_0, \cdots, T_{n_i}\}/(T_0 \cdots T_n - a_i)$ with $a \in A$ and $n \geq 0$. It is said to be* nondegenerate *if one can choose X', Y' and (B_i, a_i) such that $\{x \in (\operatorname{Spf}(A)_\eta) | a_i(x) = 0\}$ is nowhere dense.*

(ii) ϕ is said to be polystable *if there exists a surjective étale morphism $\mathfrak{Y}' \to \mathfrak{Y}$ such that $\mathfrak{Y}' \to \mathfrak{X}$ is strictly polystable. It is said to be* nondegenerate *if one can choose $\mathfrak{Y}'$ such that $\mathfrak{Y}' \to \mathfrak{X}$ is nondegenerate.*

Then a (*nondegenerate*) *polystable fibration* of length l over $\mathfrak{S}$ is a sequence of (nondegenerate) polystable morphisms $\underline{\mathfrak{X}} = (\mathfrak{X}_l \to \cdots \to \mathfrak{X}_1 \to \mathfrak{S})$.

Berkovich defines *polysimplicial sets* in [3, section 3] as follows.

For an integer n, denote $[n] = \{0, 1, \cdots, n\}$.

For a tuple $\mathbf{n} = (n_0, \cdots, n_p)$ with either $p = n_0 = 0$ are $n_i \geq 1$ for all i, let $[n]$ denote the set $[n_0] \times \cdots \times [n_p]$ and $w(\mathbf{n})$ will denote the number p.

Berkovich defines a category $\mathbf{\Lambda}$ whose objects are $[\mathbf{n}]$ and morphisms are maps $[\mathbf{m}] \to [\mathbf{n}]$ associated with triples as follows. J is a subset of $[w(\mathbf{m})]$ assumed to be empty if $[\mathbf{m}] = [0]$, f is an injective map $J \to [w(\mathbf{n})]$ and $\alpha = \{\alpha_l\}_{0 \leq l \leq p}$, where α_l is an injective map $[m_{f^{-1}(l)}] \to [n_l]$ if $l \in \mathrm{Im}(f)$, and α_l is a map $[0] \to [n_l]$ otherwise. The map $\gamma : [\mathbf{m}] \to [\mathbf{n}]$ associated with (J, f, α) takes $\mathbf{j} = (j_0, \cdots, j_{w(m)}) \in [m]$ to $\mathbf{i} = (i_0, \cdots, i_{w(\mathbf{n})})$ with $i_l = \alpha_l(j_{f^{-1}(l)})$ for $l \in \mathrm{Im}(f)$, and $i_l = \alpha_l(0)$ otherwise.

Then a polysimplicial set is a functor $\mathbf{\Lambda}^{\mathrm{op}} \to \mathrm{Set}$, they form a category denoted $\mathbf{\Lambda}^\circ\,\mathrm{Set}$.

Let Δ be the strict simplicial category of integers with increasing maps. Strict simplicial sets are simple examples of polysimplicial sets by extending to $\Delta^\circ\,\mathrm{Set} \to \mathbf{\Lambda}^\circ\,\mathrm{Set}$ the functor $\Delta \to \mathbf{\Lambda}$, that maps $[n]$ to itself, such that it commutes with direct limits.

Berkovich then considers a functor $\Sigma : \mathbf{\Lambda} \to \mathcal{K}e$ (where $\mathcal{K}e$ is the category of Kelley spaces, *i.e.* topological spaces X such that a subset of X is closed whenever its intersection with any compact subset of X is closed) that takes $[\mathbf{n}]$ to $\Sigma_{\mathbf{n}} = \{(u_{il})_{0 \leq i \leq p, 0 \leq l \leq n_i} \in [0,1]^{[\mathbf{n}]} | \sum_l u_{il} = 1\}$, and takes a map γ associated to (J, f, α) to $\Sigma(\gamma)$ that maps $\mathbf{u} = (u_{jk})$ to $\mathbf{u}' = (u'_{il})$ defined as follows: if $[\mathbf{m}] \neq [0]$ and $i \notin \mathrm{Im}(f)$ or $[\mathbf{m}] = [0]$ then $u'_{il} = 1$ for $l = \alpha_i(0)$ and $u'_{il} = 0$ otherwise; if $[\mathbf{m}] \neq [0]$ and $i \in \mathrm{Im}(f)$, then then $u'_{il} = u_{f^{-1}(i), \alpha_i^{-1}(l)}$ for $l \in \mathrm{Im}(\alpha_i)$ and $u'_{il} = 0$ otherwise. Berkovich thus gets a functor, the *geometric realization*, $|\;| : \mathbf{\Lambda}^\circ\,\mathrm{Set} \to \mathcal{K}e$ by extending Σ such that it commutes with direct limits.

Berkovich attached to a polystable fibration over $\underline{\mathfrak{X}} = (\mathfrak{X}_l \to \mathfrak{X}_{l-1} \to \cdots \to \mathrm{Spf}(k^\circ))$ a polysimplicial set $\mathbf{C}_l(\mathfrak{X}_s)$ (which only depends on the special fiber) and a subset of the generic fiber $\mathfrak{X}_{l,\eta}$ (this is a Berkovich space) of $\mathfrak{X}_l$ named the skeleton of $\underline{\mathfrak{X}}$ and denoted by $S(\underline{\mathfrak{X}})$ which is canonically homeomorphic to $|\mathbf{C}_l(\mathfrak{X}_s)|$ (see [3, th. 8.2]), such that $\mathfrak{X}_{l,\eta}$ retracts by a proper strong deformation onto $S(\underline{\mathfrak{X}})$.

In fact, when $\underline{\mathfrak{X}}$ is non degenerate (for example generically smooth, we will in fact only use the results of Berkovich to such polystable fibrations), the skeleton of $\underline{\mathfrak{X}}$ only depends on $\mathfrak{X}_l$ (such a formal scheme that fits into a polystable fibration will be called a *pluristable morphism*, and we will note $S(\mathfrak{X}_l)$ this skeleton) according to [4, prop. 4.3.1.(ii)]. In this case [4, prop. 4.3.1.(ii)] give a simple explicit description of $S(\mathfrak{X}_l)$. For any $x, y \in \mathfrak{X}_{l,\eta}$, we note $x \preceq y$ if for every $\mathfrak{X}' \to \mathfrak{X}_l$ and every point x' over x, there exists y' over y such that for every $f \in O(\mathfrak{X}_\eta)$, $|f(x')| \leq |f(y')|$ ($\preceq$ is a partial order on $\mathfrak{X}_{l,\eta}$). Then $S(\mathfrak{X}_l)$ is just the set of maximal points of $\mathfrak{X}_{l,\eta}$ for $\preceq$.

We will not give in full detail the construction of $\mathbf{C}_l(\mathfrak{X}_s)$ (we will not need it). Rather, we will give some examples for a polystable fibration of length 1

(they are all easily deduced from the construction explained in [3, section 3]).
If $\mathfrak{X}$ is just $\mathrm{Spf}(B_0) \times_{\mathrm{Spf}\, k^\circ} \cdots \times_{\mathrm{Spf}\, k^\circ} \mathrm{Spf}(B_{p+1})$ where each B_i is of the form $k^\circ\{T_0, \cdots, T_{n_i}\}/(T_0 \cdots T_{n_i} - a_i)$ with $a_i \in k^\circ$, $|a_i| < 1$ for $i < p$ and $|a_{p+1}| = 1$, then $\mathbf{C}_1(\mathfrak{X} \to \mathrm{Spf}(k^\circ))$ is just $[\mathbf{n}]$ with $\mathbf{n} = (n_0, \cdots, \mathbf{n}_p)$.
If $\mathfrak{X}$ is just a semistable formal scheme over $\mathrm{Spf}(k^\circ)$, then $\mathbf{C}_1(\mathfrak{X} \to \mathrm{Spf}(k^\circ))$ is a simplicial set.
For example, if $\mathcal{X}$ is a semistable model of a complete curve (then it is clearly polystable over k°), the polysimplicial set defined by Berkovich is just the graph of the stable reduction defined in the previous section (a graph can be considered as a simplicial set of dimension 1, and thus as a polysimplicial set).
If $(\overline{\mathcal{X}}, \mathcal{D})$ is a semistable model of a curve (not necessarily complete), the graph of $(\overline{\mathcal{X}})$ is the semigraph of $\mathcal{X} = \overline{\mathcal{X}} \backslash \mathcal{D}$ in which one deleted all the edges with only one branch. The retraction of $(\overline{\mathcal{X}}_\eta)^{\mathrm{an}}$ to the geometric realization of this graph restricts to a retraction of $(\mathcal{X}_\eta)^{\mathrm{an}}$ thanks to [3, cor. 8.4].

The retraction to $S(\underline{\mathfrak{X}})$ commutes with étale morphisms:

Theorem 3.2 ([3, th. 8.1]) *One can construct for every polystable deformation* $\mathfrak{X} = (\mathfrak{X}_l \xrightarrow{f_{l-1}} \cdots \xrightarrow{f_1} \mathfrak{X}_1 \to \mathrm{Spf}(k^\circ))$ *a proper strong deformation retraction* $\Phi^l : \mathfrak{X}_{l,\eta} \times [0, l] \to \mathfrak{X}_{l,\eta}$ *of* $\mathfrak{X}_{l,\eta}$ *onto the skeleton* $S(\underline{\mathfrak{X}})$ *of* $\underline{\mathfrak{X}}$ *such that :*

(i) $S(\underline{\mathfrak{X}}) = \bigcup_{x \in S(\underline{\mathfrak{X}}_{l-1})} S(\mathfrak{X}_{l,x})$*, where* $\underline{\mathfrak{X}}_{l-1} = (\mathfrak{X}_{l-1} \to \cdots \to \mathrm{Spf}(k^\circ))$*, and* $f_{l-1,\eta}(x_t) = f_{l-1,\eta}(x)_{t-1}$ *for every* $t \in [1, l]$*;*

(ii) $(x_t)_{t'} = x_{\max(t,t')}$*;*

(iii) $x \leqslant x_t$*;*

(iv) for every x *and* $i \leqslant l-1$*, there is* $t' \in [i, i+1]$ *such that* $x_t = x_i$ *if* $t \in [i, t']$ *and* $t \mapsto x_t$ *is injective over* $[t', i+1]$*;*

(v) for every t*,* $\pi(x_t)$ *is contained in the same stratum of* $\mathfrak{X}_{l,s}$ *as* $\pi(x)$*; moreover* $\pi(x_l)$ *is the generic point of this stratum and* $\pi(x_i) = \pi(x_t)$ *for every* $0 \leqslant i \leqslant l-1$ *and every* $t \in [i, i+1]$*;*

(vi) if $\phi : \mathfrak{Y} \to \mathfrak{X}$ *is a morphism of fibrations in* $\mathcal{P}stf_l^{\acute{e}t}$*, one has* $\phi_{l,\eta}(y_t) = \phi_{l,\eta}(y)_t$ *for every* $y \in \mathfrak{Y}_{l,\eta}$*.*

Berkovich deduces from the functoriality of the retraction with respect to étale morphisms the following corollary:

Corollary 3.3 ([3, cor 8.5]) *Let* $\underline{\mathfrak{X}}$ *be a polystable fibration over* k'° *(where* k' *is a finite normal extension of* k*) with a normal generic fiber* $\mathfrak{X}_{l,\eta}$*. Suppose we are given an action of a finite group* G *on* $\underline{\mathfrak{X}}$ *over* k° *and a Zariski open dense subset* U *of* $\mathfrak{X}_{l,\eta}$*. Then there is a strong deformation retraction of the Berkovich space* $G\backslash U$ *to a closed subset homeomorphic to* $G\backslash|\mathbf{C}^l(\underline{\mathfrak{X}})|$*.*

More precisely, in this corollary, the closed subset in question is the image of $S(\underline{\mathfrak{X}})$ (which is G-equivariant and contained in U) by $U \to G\backslash U$.

Recall that the skeleton is functorial for pluristable morphisms:

Proposition 3.4 ([4, prop. 4.3.2.(i)]) *If $\phi : \mathfrak{X} \to \mathfrak{Y}$ is a pluristable morphism between nondegenerate pluristable formal schemes over k°, $\phi_\eta(S(\mathfrak{X})) \subset S(\mathfrak{Y})$.*

In fact, more precisely, from the construction of S, $S(\mathfrak{Y}) = \bigcup_{x \in S(\mathfrak{X})} S(\mathfrak{Y}_x)$. Recall also that, if $\bar{k}$ is algebraically closed, the polysimplicial complex of a polystable fibration commutes with base change:

Proposition 3.5 ([3, prop. 6.10]) *If $\underline{X}$ is a polystable fibration over a field K, there exists a finite separable extension K' of K such that for any bigger field K'', $\mathbf{C}(\underline{X} \otimes K'') \to \mathbf{C}(\underline{X} \otimes K')$ is an isomorphism.*

In order to use the previous description of the Berkovich space of a scheme with a model over k° which admits a polystable fibration for understanding the topology of a smooth scheme over k, we will need de Jong's result about the existence of alterations by such pluristable schemes over k°.
More precisely we will use the following consequence of de Jong's theory given by Berkovich (as we will work over valuation fields of characteristic 0, we give here only a restricted version in this case):

Lemma 3.6 ([3, lem. 9.2]) *Let A be a Henselian valuation ring with fraction field of characteristic 0, let X be an integral scheme proper flat and finite presentation over A, with an irreducible generic fiber of dimension l. Then there is:*

(a) a finite Galois extension of the fraction field of A, with ring of integers A',

(b) a polystable fibration $\underline{X}' = (X'_l \to \cdots \to X'_0 = \operatorname{Spec} A')$, where every morphism is projective of dimension 1 with smooth geometrically irreducible generic fibres,

(c) an action of a finite group G on $\underline{X}'$ over A,

(d) a commutative diagram

$$\begin{tikzcd}
X'_l \arrow[r, "\phi"] \arrow[d] & X \arrow[d] \\
\operatorname{Spec}(A') \arrow[r] & \operatorname{Spec}(A)
\end{tikzcd}$$

where ϕ is a dominant G-equivariant morphism such that the generic fiber is generically étale with Galois group G.

3.2 Invariance of π_1^{temp} by base change of algebraically closed fields

Let X be a smooth algebraic variety over an algebraically closed complete nonarchimedean field K of characteristic 0.
Let $X_i \to X$ be a finite étale connected Galois covering and U_0 a dense affine Zariski open subset of X_i, and let us embed (*e.g.* by [3, lem 9.4]) U_0 in a scheme $\overline{X_i}$ which is proper , of finite presentation and flat over O_K.
Then there is by lemma (3.6) a generically smooth polystable fibration X'_i over

O_K endowed with an action of a group G such that (X'_i, G) is a Galois alteration of $\overline{X_i}$.
Let U be a dense Zariski open subset of $\overline{X_i}$ included in U_0 (and therefore in X_i) such that $U' \to U$ is finite (where U' is the pullback of U in X'_i).
Then U^{an} retracts by strong deformation on $G\backslash S(X'_{i,s})$ by corollary (3.3).

Let K'/K an algebraically closed extension of complete valued fields and let x' a geometric point with image.
$X_{i,K'} \to X_{K'}$ is also a finite étale connected Galois covering (by [10, cor X.1.8]), $X'_{i,0_{K'}}$ is also a polystable fibration endowed with an action of G and $(X'_{i,O_{K'}}$ is also a Galois alteration of $\overline{X}_{i,K'}$, finite over $U_{K'}$.
Thus, as in the previous case, $U^{\mathrm{an}}_{K'}$ retracts on the closed subset $G\backslash S(X'_{i,K'})$ (and the natural morphism $U^{\mathrm{an}}_{K'} \to U^{\mathrm{an}}$ maps $S(X'_{i,K',s})$ to $S(X'_{i,s})$.
But $\mathbf{C}(X'_{i,K',s}) \to \mathbf{C}(X'_{i,s})$ is an isomorphism according to proposition (3.5). The morphism $U^{\mathrm{an}}_{K'} \to U^{\mathrm{an}}$ is thus a homotopy equivalence.
One has the following 2-commutative diagram :

$$\begin{tikzcd}
\mathrm{Cov}^{\mathrm{top}}(U_{K'}) & \mathrm{Cov}^{\mathrm{top}}(U) \arrow[l] \\
\mathrm{Cov}^{\mathrm{top}}(X_{i,K'}) \arrow[u] & \mathrm{Cov}^{\mathrm{top}}(X_i) \arrow[l] \arrow[u]
\end{tikzcd} .$$

The vertical arrows are equivalences by proposition (1.4), and the top arrow is also an equivalence by what has just been shown.
Thus, if x'_i is a geometric point of $X_{i,K'}$ with image x_i in X_i, $\pi_1^{\mathrm{top}}(X_{i,K'}, x'_i) \to \pi_1^{\mathrm{top}}(X_i, x_i)$ is an isomorphism.
If X_i^{∞} is the universal topological covering of X_i and $X^{\infty}_{i,K'}$ is the universal topological covering of $X_{i,K'}$, one has $X^{\infty}_{i,K'} = (X_i^{\infty})_{K'}$.
Therefore $\mathrm{Gal}(X^{\infty}_{i,K'}/X_{K'}) = \mathrm{Gal}(X^{\infty}_i/X)$.
By taking the projective limit over a cofinal projective system of geometrically pointed (the points being defined in a big enough valuation field Ω) Galois coverings $((X_i)_i, x_i)_{i\in\mathbf{N}}$ $((X_{i,K'}, x'_i)_{i\in\mathbf{N}}$, where x'_i is some point of $X_{i,K'}$ over x_i, is then also a cofinal projective system of Galois covering of $X_{K'}$ by [10, lecture XIII]), one gets :

Proposition 3.7 *The morphism* $\pi_1^{\mathrm{temp}}(X_{K'}, x') \to \pi_1^{\mathrm{temp}}(X, x)$ *is an isomorphism.*

3.3 Products and tempered fundamental group

Let Y be another smooth algebraic variety over K, $Y_j \to Y$ a finite étale connected Galois covering, $\overline{Y_j}$ a scheme which is proper, of finite presentation and flat over O_K, in which is embedded a dense affine open subset V_0 of Y_j, and let $(Y'_j, H) \to \overline{Y_j}$ a Galois alteration and $Y'_j \to O_K$ a polystable fibration.
Let also $V \subset Y_j$ be a dense Zariski open subset, included in V_0 such that $V' \to V$ is finite.
Then, as for X, V^{an} retracts by strong deformation over $H\backslash|\mathbf{C}(Y'_{j,s})|$.

One get the same results for $X_i \times Y_j$:
$X'_i \times Y'_j \to X_i \times Y_j$ is a Galois alteration of group $G \times H$, and $U \times V$ retracts

on $(G \times H)\backslash S(X_i \times Y_j)$.
But the pluristable maps $X_i \times Y_j \to X_i$ and $X_i \times Y_j \to Y_j$ map $S(X_i \times Y_j)$ into $S(X_i)$ and $S(X_j)$ respectively. Whence a map $f : S(X_i \times Y_j) \to S(X_i) \times S(Y_j)$ (and it is compatible with the action of $G \times H$). But as

$$S(X_i \times Y_j) = \bigcup_{x \in S(X_i)} S((X_i \times Y_j)_x) = \bigcup_{x \in S(X_i)} S(Y_j \otimes \mathcal{H}(x))$$

and $S(Y_j \otimes \mathcal{H}(x)) \to S(Y_j)$ is an isomorphism according to proposition (3.5), f is bijective. But $S(X_i \times Y_j)$ is compact, and thus f is an isomorphism.
Thus $(G \times H)\backslash S(X_i \times Y_j) \to G\backslash S(X_i) \times H\backslash S(Y_j)$ is an isomorphism.
Thus $(U \times V)^{\mathrm{an}} \to U^{\mathrm{an}} \times V^{\mathrm{an}}$ is a homotopy equivalence (the product on the right is the product of topological spaces), and by applying proposition (1.4) to $U \subset X_i$, $V \subset Y_j$ and $U \times V \subset X_i \times Y_j$, one gets that

$$\pi_1^{\mathrm{top}}(X_i \times Y_j, (x_i, y_j)) \to \pi_1^{\mathrm{top}}(X_i, x_i) \times \pi_1^{\mathrm{top}}(Y_j, y_j)$$

is an isomorphism.
So $(X_i \times Y_j)^\infty = X_i^\infty \times Y_j^\infty$ and

$$\mathrm{Gal}((X_i \times Y_j)^\infty/(X \times Y)) = \mathrm{Gal}(X_i^\infty/X) \times \mathrm{Gal}(Y_j^\infty/Y).$$

If $(X_i, x_i)_i$ and $(Y_j, y_j)_j$ are cofinal projective systems of connected geometrically pointed Galois coverings of X and Y, $(X_i \times Y_j, (x_i, y_j))_{(i,j)}$ is a cofinal projective system of connected Galois coverings of $X \times Y$ by [10, lecture XIII].
By taking the projective limit over (i, j) in the previous isomorphism, one gets:

Proposition 3.8 $\pi_1^{\mathrm{temp}}(X \times Y, (x, y)) \to \pi_1^{\mathrm{temp}}(X, x) \times \pi_1^{\mathrm{temp}}(Y, y)$ *is an isomorphism.*

4 Metric structure of the graph of the stable model of a curve

The main goal of this section is to prove that the metric structure of the graph of the stable model (or equivalently, the graph structure of the skeleton of the curve considered as a Berkovich space, as recalled in section 3.1) of a Mumford curve can be recovered from the tempered fundamental group of this curve. We will only consider the case of mixed characteristic (in the case of equal characteristic 0, this is surely false, as the whole tempered fundamental group can be recovered only from the graph of groups associated with the stable reduction of the curve according to [13, ex. 3.10]).

Definition 4.1 *A* metric structure *over a graph* $\mathbb{G}$ *is a function*

$$d : \{\textit{edges of } \mathbb{G}\} \to \mathbf{R}^{>0}.$$

For any e, $d(e)$ *is called the* length *of the edge* e *with respect to the metric structure* d.
A graph endowed with a metric structure is called a metric graph.

For a curve X over an algebraically closed complete nonarchimedean field k with a semistable model $\mathcal{X}$, let e be an edge of the graph of this semistable reduction (that is a node of $\mathcal{X}_s$). Then etale locally at this node, $\mathcal{X}$ is étale over $k^\circ[X_0, X_1]/(X_0X_1 - a)$ with $a \in k^\circ$. According to [14, cor. 2.2.18], $|a|$ does not depend of any choice. Then we will denote $d(e) = -\log_p(|a|)$, which defines a natural metric structure on the graph of the stable reduction of X.
For example, if $\mathcal{X}$ is the stable model of $\mathbf{P}^1\backslash\{0, 1, \infty, \lambda\}$ with $|\lambda| < 1$ then the graph of $\mathcal{X}$ has a single edge of length $-\log_p(|\lambda|)$.

We already know, thanks to [13, ex. 3.10] that one can recover the graph of the stable model from the tempered fundamental group. In fact we will deduce from Mochizuki's study that one can recover, for every finite index open subgroup of the tempered fundamental group and every vertex of the skeleton of the curve, whether the corresponding covering of the curve is split over this vertex (an étale covering $X' \to X$ of manifolds will be said to be *split* over a point $x \in X$ if, for every $x' \in X'$ over x, $\mathcal{H}(x) \to \mathcal{H}(x')$ is an isomorphism; if $X' \to X$ is a covering of order n, then it is split over x if and only if the fiber of x has cardinality n, if and only if, locally in a neighborhood of x, $X' \to X$ pulls back to a topological covering thanks to [2, III.1.2.1]).
This suggests to consider how finite étale coverings of a Mumford curve are split over vertex points. No one knows, as far as I know, how to determine in general if a covering is split over some point of the Berkovich space. However, studying simple coverings may be enough to see that the metric structure of the skeleton must play a role in the structure of the tempered fundamental group.
More precisely,

Lemma 4.2 *The covering* $\mathbf{G}_m \overset{z\mapsto z^{p^e}}{\to} \mathbf{G}_m$ *is split over a Berkovich point* $B(1,r)$ *corresponding to the ball of center 1 and ray* r *with* $r < 1$, *if and only if* $r < p^{-e-\frac{1}{p-1}}$.
More precisely, $B(1,r)$ *has* p^i *preimage if:*

- $r \in]p^{-\frac{p}{p-1}}, 1]$, *if* $i = 0$*;*
- $r \in]p^{-i-\frac{p}{p-1}}, p^{-i-\frac{1}{p-1}}]$, *if* $1 \leq i \leq e-1$*;*
- $r \in [0, p^{-e-\frac{1}{p-1}}]$, *if* $i = e$.

Proof : Let $g : z \mapsto z^p$, and let us calculate $g(B(h', r))$ with $|h'| = 1$ and $r < 1$. Let $f'_h : z \mapsto (z + h')^p - h'^p = \sum_{i=1}^p a_i z^i$ with $|a_p| = 1$ and $|a_i| = p^{-1}$ if $1 \leq i \leq p-1$.
Then $g(B(h', r)) = B(h'^p, r')$ with

$$r' = \max_{|z|<r} |f(z)| = \max |a_i| r^i = \max\{r^p, rp^{-1}\} = \begin{cases} rp^{-1} & \text{if } r \leq p^{-\frac{1}{p-1}} \\ r^p & \text{if } r \geq p^{-\frac{1}{p-1}} \end{cases}$$

Moreover let h be of norm 1, let $h^{1/p}$ be a pth root of h, and let $r' < 1$. Then $|h'^p - h| = \prod_{\zeta\in\mu_p} |h' - \zeta h^{1/p}| \leq r'$ implies that there exists $\zeta_0 \in \mu_p$ such that $|h' - \zeta_0 h^{1/p}| \leq r'^{1/p}$ (*i.e.* $B(h', r'^{1/p}) = B(h^{1/p}, r'^{1/p})$).
Suppose now $|\zeta - \zeta'| = p^{-\frac{1}{p-1}}$. Since $|\zeta - \zeta_0| = p^{-\frac{1}{p-1}}$ if $\zeta \in \mu_p\backslash\{\zeta_0\}$, $|h' -$

$h^{1/p}\zeta| = p^{-\frac{1}{p-1}}$ if $\zeta \in \mu_p \backslash \{\zeta_0\}$ and thus $|h' - h^{1/p}\zeta_0| = |h'^p - h| / \prod_{\zeta \in \mu_p \backslash \{\zeta_0\}} |h' - h^{1/p}\zeta| \leq pr'$ (*i.e.* $B(h', pr') = B(\zeta_0 h^{1/p}, pr')$). Thus

$$g^{-1}(B(h,r')) = \left\{ \begin{array}{ll} \{B(\zeta h^{1/p}, pr')\}_{\zeta \in \mu_p} & \text{if } r' \leq p^{-\frac{p}{p-1}} \\ \{B(\zeta h^{1/p}, r'^{1/p})\}_{\zeta \in \mu_p} & \text{if } r' \geq p^{-\frac{p}{p-1}} \end{array} \right.$$

Since $|\zeta - \zeta'| = p^{-\frac{1}{p-1}}$ if $\zeta \neq \zeta' \in \mu_p$, one gets that $g^{-1}(B(h,r'))$ has a single element if $r' \geq p^{-\frac{p}{p-1}}$ and p otherwise. Thus ones gets the wanted result when $e = 1$.
In the general case, one uses induction on e by decomposing $z \mapsto z^{p^e}$ in $z \mapsto z^{p^{e-1}} \mapsto z^{p^e}$. □

Thanks to this, we will first study the case of the projective line minus some points and, by cutting and pasting this kind of covering, the case of a punctured elliptic curve.
For the more general case of a Mumford curve X, we will also study the structure of $\mathbf{Z}/p^e\mathbf{Z}$-torsors. The theory of theta functions as can be found in [16] and [15] tells us that the pullback to the universal topological covering Ω of such a torsor is in fact the pullback of $\mathbf{G}_m \overset{z\mapsto z^{p^e}}{\to} \mathbf{G}_m$ along some theta function $\Omega \to \mathbf{G}_m$, which in turn corresponds to some equivariant current over the tree $\mathbb{T}(\Omega)$ of Ω. Therefore, we will begin our study by proving that if two currents coincide over a sufficiently big part of $\mathbb{T}(\Omega)$, the quotient of the two corresponding invertible functions is nearly constant over some smaller part of $\mathbb{T}(\Omega)$ and thus the two corresponding $\mathbf{Z}/p^e\mathbf{Z}$-torsors are split over the same vertices in this part of $\mathbb{T}(\Omega)$. Thus, we will consider some currents which, over some "big" part of Ω, coincides with the one corresponding to a homography invertible over ω and which is equivariant under some subgroup of finite index of $\mathrm{Gal}(\Omega/X)$. Then we consider the corresponding $\mathbf{Z}/p^e\mathbf{Z}$-torsor over some finite étale covering of X which will behave like $\mathbf{G}_m \overset{z\mapsto z^{p^e}}{\to} \mathbf{G}_m$ over some part of $\mathbb{T}(\Omega)$.
We will deduce from this that the length of every loop of every finite topological covering of the graph of the stable model of X can be recovered from the tempered fundamental group. A final combinatorial consideration will give us what we wanted.

4.1 Preliminaries

Let K be a complete nonarchimedean field of characteristic 0, K° its integral ring, k its residual field, assumed of characteristic $p > 0$, and let $\overline{K}$ be the completion of the algebraic closure of K (with integral ring $\overline{K}^\circ$).
From now on, we assume that we have chosen a compatible system of roots of 1 in $\overline{K}$, so that we may identify μ_n and $\mathbf{Z}/n\mathbf{Z}$. Thus we will often talk of $\mathbf{Z}/n\mathbf{Z}$-torsors over a curve over $\overline{K}$ when we should better talk of μ_n-torsors.
Let X_K a smooth curve of type (g,n) over K and let $Y_K \to X_K$ a geometrically connected finite étale covering of X_K.

Let $\Pi = \pi_1^{\mathrm{temp}}(X_{\overline{K}})$ and $H \subset \Pi$ the open subgroup of finite index corresponding to Y_K.

By changing K by a finite extension, one may assume that X_K and Y_K have a stable model over K°, denoted X and Y.
One then have a unique morphism $\psi : Y \to X$ extending $Y_K \to X_K$.

By theorem (1.7), one may rebuild from the topological group Π the semigraph of groups $\mathcal{G}$ of $X_{\overline{k}}$. Likewise, one may rebuild from H the semigraph of groups $\mathcal{H}$ of $Y_{\overline{k}}$.
We are now interested in reconstructing from $\pi_1^{\text{temp}}(Y_K) \to \pi_1^{\text{temp}}(X_K)$ what data can be recovered of the preimage of the cusps and vertices of the skeleton of X_K.

4.1.1 Cusps

Let y be a cusp of $Y_{\overline{k}}$ (corresponding to a cusp y' of $Y_{\overline{K}}$) and let x be its image in $X_{\overline{k}}$ (corresponding to a cusp x' of $X_{\overline{K}}$).
Let $I \subset H^{(p')}$ be an inertia group of y. The image of I in $\Pi^{(p')}$ is an open subgroup (and thus it is nontrivial) of an inertia group of x. Since the intersection of the inertia group of two different cusps is $\{1\}$, the image of I is not contained in any other inertia group of a cusp of $X_{\overline{k}}$, thus x is characterized by the morphism $H \to \Pi$ as being the only cusp of $X_{\overline{K}}$ such that the inertia groups of y map by $H \to \Pi$ to inertia groups of x.
Thus one may rebuild from $H \subset \Pi$ the map from the cusps of $Y_{\overline{k}}$ to the cusps of $X_{\overline{k}}$. In particular, one may decide whether the morphism $Y \to X$ is ramified at x'.

4.1.2 Vertices of the skeleton

If X_0 is an irreducible component of $X_{\overline{k}}$, let Y_0 be an irreducible component of $Y_{\overline{k}}$ which maps surjectively to X_0. Then the morphism between the components of the graphs of groups $\Pi_{Y_0}^{(p')} \to \Pi_{X_0}^{(p')}$ is open (in particular, its image is non commutative) since it embeds in the following commutative diagram

$$\begin{tikzcd}
\operatorname{Gal}\left(\overline{K(Y_0)}/K(Y_0)\right) \arrow[r, hook] \arrow[d, two heads] & \operatorname{Gal}\left(\overline{K(X_0)}/K(Y_0)\right) \arrow[d, two heads] \\
\Pi_{Y_0}^{(p')} \arrow[r] & \Pi_{X_0}^{(p')}
\end{tikzcd},$$

where the upper arrow is an open embedding and the vertical arrows are projections.
Since $\Pi_{X_0}^{(p')} \to \Pi^{(p')}$ (defined up to conjugation) is injective, the image of $\Pi_{Y_0}^{(p')}$ in $\Pi^{(p')}$ (defined up to conjugation) is non commutative, and thus $\Pi_{X_0}^{(p')}$ is the only verticial subgroup of $\Pi^{(p')}$ which contains the image of $\Pi_{Y_0}^{(p')}$.
Moreover, if Y_0 is an irreducible component of $Y_{\overline{k}}$ which does not map surjectively onto an irreducible component of $X_{\overline{k}}$, the image of $\Pi_{Y_0}^{(p')}$ in $\Pi^{(p')}$ is commutative, so the embedding $H \to \Pi$ decides which components of $Y_{\overline{k}}$ maps surjectively onto which components of $X_{\overline{k}}$.

In particular, one may rebuild from $H \to \pi$ the cardinal of the preimage by $Y_{\overline{k}}$ of the generic points of $X_{\overline{k}}$.
Equivalently, if x_0 is a vertex of the skeleton of $X^{\mathrm{an}}_{\overline{K}}$, one may rebuild the cardinal of its preimage in $Y^{\mathrm{an}}_{\overline{K}}$.
Indeed, let us write π for the continuous map from the generic analytic fiber to the special fibre. $x_1 = \pi(x_0)$ is the generic point of an irreducible component of $X_{\overline{k}}$, every preimage y_1 of x_1 by ψ is a generic point of an irreducible component of $Y_{\overline{k}}$, $\pi^{-1}(y_1)$ is reduced to a single element by [3, cor. 1.7], which must map to x_0 because $\pi^{-1}(x_1) = \{x_0\}$ by [3, cor. 1.7]. Thus $\psi^{\mathrm{an},-1}(x_0)$ is in natural bijection with $\psi^{-1}(x_1)$.

4.2 Case of $\mathbf{P}^1\backslash\{z_1, \dots, z_n\}$

Let $\Box = \alpha, \beta$.
Let $z_{\Box,1}, \cdots, z_{\Box,n} \in \mathbf{Q}_p^{\mathrm{nr}}$, with $n \geqslant 4$.
Let us write $X_\Box = \mathbf{P}^1\backslash\{z_{\Box,1}, \dots, z_{\Box,n}\}$.
Let $\Pi_\Box = \pi_1^{\mathrm{temp}}(\mathbf{P}^1\backslash\{z_{\Box,1}, \dots, z_{\Box,n}\})$. We already know that an isomorphism $\phi : \Pi_\alpha \simeq \Pi_\beta$ induces an isomorphism between the semigraphs of the stable reductions of $\mathbf{P}^1\backslash\{z_{\alpha,1}, \dots, z_{\alpha,n}\}$. After reordering the $z_{\beta,i}$, we may assume that this morphism of semigraphs identifies the inertia subgroup (defined up to conjugation) of the cusp $z_{\alpha,i}$ with the inertia subgroup of the cusp $z_{\beta,i}$.

Theorem 4.3 *The isomorphism of graphs thus defined by ϕ between the skeletons of $(\mathbf{P}^1\backslash\{z_{\alpha,1}, \dots, z_{\alpha,n}\})^{\mathrm{an}}$ and $(\mathbf{P}^1\backslash\{z_{\beta,1}, \dots, z_{\beta,n}\})^{\mathrm{an}}$ preserves lengths of edges (*i.e. *it induces an isomorphism of metric graphs).*

Equivalently, for every (i_1, i_2, i_3, i_4), one gets the equality of cross-ratios' norms:

$$|(z_{\alpha,i_1}, z_{\alpha,i_2}, z_{\alpha,i_3}, z_{\alpha,i_4})| = |(z_{\beta,i_1}, z_{\beta,i_2}, z_{\beta,i_3}, z_{\beta,i_4})|.$$

In fact, we will be able to prove this result without assuming $z_{\Box,1}, \cdots, z_{\Box,n} \in \mathbf{Q}_p^{\mathrm{nr}}$ after studying the case of an elliptic curve (this will give the result for $p \neq 2$) and without any assumption after studying the case of a Mumford curve.

Proof : According to section 4.1.1, if H_α and $H_\beta = \phi(H_\alpha)$ are finite index open subgroups corresponding to each other by ϕ, we know that the corresponding covering of X_α is ramified outside $\{z_{\alpha,i_1}, z_{\alpha,i_2}, z_{\alpha,i_3}, z_{\alpha,i_4}\}$ if and only if the corresponding covering of X_β is ramified outside $\{z_{\beta,i_1}, z_{\beta,i_2}, z_{\beta,i_3}, z_{\beta,i_4}\}$.
Thus ϕ induces an isomorphism between the $\pi_1^{\mathrm{temp}}(\mathbf{P}^1\backslash\{z_{\Box,i_1}, z_{\Box,i_2}, z_{\Box,i_3}, z_{\Box,i_4}\})$ (it is the projective limit of the discrete quotients $G_\Box$ of $\Pi_\Box$ which have a finite quotient $G_{1,\Box}$ corresponding to a covering of $\mathbf{P}^1$ unramified outside $\{z_{\Box,i_1}, z_{\Box,i_2}, z_{\Box,i_3}, z_{\Box,i_4}\}$ such that $\mathrm{Ker}(G_\Box \to G_{1,\Box})$ is a free group (or, equivalently here, is torsion free)).

We thus may assume that $n = 4$. We may also assume, after a base change of an homography, that the cusps $0, 1, \infty$ are $\lambda_\Box$. Moreover, we may assume that $X_\Box$ does not have good reduction (in which case there is nothing to prove), and thus that $v_p(\lambda_\Box - 1) > 0$ (it is an integer since $\lambda \in \mathbf{Q}_p^{\mathrm{nr}}$) after permuting $0, 1$ and ∞ by another homography. We now have to prove that $v_p(\lambda_\alpha - 1) = v_p(\lambda_\beta - 1)$. Assume, ab absurdo, that $v_p(\lambda_\alpha - 1) < v_p(\lambda_\beta - 1)$. Let $e := v_p(\lambda_\beta - 1) - 1 \geqslant v_p(\lambda_{alpha} - 1)$ and let $H_\Box$ be the subgroup of $\Pi_\Box$ of index p^e corresponding to

the unique connected covering $Y_\Box \to X_\Box$ of degree p^e unramified outside 0 and ∞ (this is the morphism $z \mapsto z^{p^e}$ from $\mathbf{P}^1$ to $\mathbf{P}^1$). One has $\phi(H_\alpha) = H_\beta$ by section 4.1.1.
But, according to lemma (4.2), $B(1, r)$ (the point of the Berkovich space of $\mathbf{P}^1$ corresponding to the ball of radius r and center 1) has p^{e-1} preimages if $p^{-(e+\frac{1}{p-1})} \leqslant r < p^{-(e-1+\frac{1}{p-1})}$ and has p^e preimages if $r < p^{-(e+\frac{1}{p-1})}$.
Thus if $p \neq 2$, $B(1, |\lambda_\alpha - 1|)$ has p^{e-1} preimages if Y_α and $B(1, |\lambda_\beta - 1|)$ has p^e preimages in Y_β, which contradicts the result of part 4.1.2.

In the case $p = 2$ and $e \geqslant 2$, $B(1, |\lambda_\alpha - 1|)$ has 2^{e-2} preimages in Y_α and $B(1, |\lambda_\beta - 1|)$ has 2^{e-1} preimages in Y_β, which contradicts the result of part 4.1.2.
If $e = 1$, and therefore $v_2(\lambda_\alpha) = 1$ end $v_2(\lambda_\beta) = 2$, the semigraph of the reduction of $Y_\alpha \to X_\alpha$ is (we marked the different cusps of Y_α, and $\sqrt{\lambda_\Box}$ is a square root of $\lambda_\Box$):



The semigraph of the stable reduction of Y_β is



They are not isomorphic, and thus one also gets a contradiction. $\Box$

4.3 Case of a punctured elliptic curve

Let $\Box = \alpha, \beta$
Let $X_\Box$ be two punctured Tate curves $\mathbf{C}_p^*/q_\Box^{\mathbf{Z}} - \{1\}$ with $|q_\Box| < 1$.

Let $\Pi_\square = \pi_1^{\text{temp}} X_\square)$ and let $\phi : \Pi_\alpha \simeq \Pi_\beta$ be an isomorphism.

Theorem 4.4 $|q_\alpha| = |q_\beta|$

Proof :
Let us choose integers n, l and m such that:

- n is prime to p and$n \geqslant \frac{v_p(q_\beta)v_p(q_\alpha)(p-1)}{|v_p(q_\beta)-v_p(q_\alpha)|p}$,
- $l \geqslant 1 + \frac{2np}{(p-1)v_p(q_\square)}$,
- $m \geqslant \frac{2l}{n}$.

Let $H_{0,\square} = [\Pi_\square, \Pi_\square]\Pi_\square^n$ be the preimage in $\Pi_\square$ of the image by the multiplication by n in the abelianized group of $\Pi_\square$. ϕ induces an isomorphism $H_{0,\alpha} \to H_{0,\beta}$.
The covering $Y_{0,\square}$ of $X_\square$ corresponding to $H_\square$ is the multiplication $\overline{X}_\square \overset{\times n}{\to} \overline{X}_\square$ by n on the elliptic curve $\overline{X}_\square$.

Let now $H_{1,\square}$ be the subgroup of $H_{0,\square}$ corresponding to the unique connected topological covering $Y_{1,\square}$ of degree n of $Y_{0,\square}$.
$Y_{1,sq} \simeq \mathbf{C}_p^*/q^{m\mathbf{Z}} - \{q^{\frac{a}{n}}\zeta^b\}_{(a,b)\in\mathbf{Z}^2}$ où $q^{\frac{1}{n}}$ is an n-th root of q and ζ is an n-th root of 1.
The semigraph of the stable reduction of $Y_{1,\square}$ has mn vertices joined in circle (the distance of such two successive vertices is $v_p(q_\square)/n$) and n cusps abuts to each vertex.
ϕ induces an isomorphism $H_{1,\alpha} \simeq H_{1,\beta}$ which itself induces an isomorphism between the semigraphs of the stable reduction of $Y_{1,\square}$. Let us number from 0 to $mn-1$ the vertices of the graphs by following the circle compatibly with the isomorphism induced by ϕ (let us write $x_{\square,0}, \cdots, x_{\square,mn-1}$ for the corresponding vertices of the skeleton of $Y_{1,\square}$).
Let $z_{\alpha,0}$ and $z_{\alpha,,l}$ be two cusps of $Y_{1,\alpha}$ abutting to the vertices of the graph numbered 0 and l respectively. Let $z_{\beta,0}$ and $z_{\beta,1}$ be the corresponding cusps of $Y_{1,\beta}$.

Let us now focus on $\mathbf{F}_p$-torsors over $\overline{Y}_{1,\square}$ which are unramified outside $z_{\square,0}$ and $z_{\square,l}$. They are the elements of an $\mathbf{F}_p$-vector space $V_\square$ of dimension 3.
Recall that we chose m, n and l so that $\frac{l-1}{n}v_p(q_\square) > \frac{2p}{p-1}$ and $\frac{mn-l}{n}v_p(q_\square) > \frac{l}{n}v_p(q_\square)$.
We will now describe a base of this $\mathbf{F}_p$-vector space.

Let $S_\square$ be the universal topological covering of $\overline{Y}_{1,\square}$, we identify it with $\mathbf{P}^1\backslash\{0,\infty\} \subset \mathbf{P}^1$. Let $s_{\square,0}$ and $s_{\square,l}$ the unique preimages in $S_\square$ of $Z_{\square,0}$ and $z_{\square,l}$ of norm 1 and $|q|^{\frac{l}{mn}}$ and let $U_{1,\square} \subset S_\square$ be the open crown $\{|q_\square|^{\frac{l-mn}{2n}} > |z| > |q_\square|^{\frac{l+mn}{2n}}\}$ and $U_{2,\square} \subset S_\square$ the open annulus $\{|q_\square|^{\frac{l}{n}}p^{-\frac{p}{p-1}} > |z| > |q_\square|^m p^{\frac{p}{p-1}}\}$. The maps fom $U_{1,\square}$ and from $U_{2,\square}$ to $\overline{Y}_{1,\square}$ still are open embeddings which, together, cover $\overline{Y}_{1,\square}$.
Let $T_{1,\square}$ the restriction to $U_{1,\square}$ of the ramified (only over $s_{\square,0}$ and $s_{\square,l}$) covering $\mathbf{P}^1 \to \mathbf{P}^1 : z \mapsto \frac{s_{\square,0}z^p+s_{\square,l}}{z^p+1}$, which is Galois of Galois group isomorphic to $\mathbf{Z}/p\mathbf{Z}$, and let us choose such an isomorphism to get a $\mathbf{Z}/p\mathbf{Z}$-torsor. Let $T_{2,\square}$

be the trivial $\mathbf{Z}/p\mathbf{Z}$-torsor over $U_{2,\Box}$ and let $T_{3,\Box} = T_{1,\Box} \coprod T_{2,\Box} \to U_{3,\Box} = U_{1,\Box} \coprod U_{2,\Box}$.
Over $U_{1,\Box} \times_{\overline{Y}_{1,\Box}} U_{2,\Box}$ (which has two connected components), $T_{1,\Box}$ is trivial. Let us choose a trivialization : one may now descend $T_{3,\Box} \to U_{3,\Box}$ into a $\mathbf{Z}/p\mathbf{Z}$-torsor $T_\Box \to \overline{Y}_{1,\Box}$, which is only ramified over $z_{\Box,0}$ and $z_{\Box,l}$.
According to lemma (4.2), $T_\Box \to \overline{Y}_{1,\Box}$ is split over x_i (with $i \in [0, mn-1]$) if and only if $|q_\Box|^{\frac{i}{n}} \in [|q_\Box|^m p^{\frac{p}{p-1}}, |q_\Box|^{\frac{l}{n}} p^{-\frac{p}{p-1}}]$, *i. e.*

$$i \in I_{1,\Box} := [l + \frac{np}{v_p(q_\Box)(p-1)}, mn - \frac{np}{v_p(q_\Box)(p-1)}].$$

There is such an integer thanks to the assumption about l, m and n because

$$\lg(I_{1,\Box}) = mn - l - 2\frac{np}{v_p(q_\Box)(p-1)} \geqslant 1$$

(where lg denotes the length of an interval).

Likewise, let $s'_{\Box,l}$ be a preimage of $z_{\Box,l}$ of norm $|q|^{\frac{l-mn}{n}}$, let $U'_{1,\Box}$ be the crown $\{|q_\Box|^{\frac{l-2mn}{2n}} > |z| > |q_\Box|^{\frac{l}{2n}}\}$, $U'_{2,\Box}$ the crown $\{p^{-\frac{p}{p-1}} > |z| > |q_\Box|^{\frac{mn-l}{n}} p^{\frac{p}{p-1}}\}$. They are open subsets of $\overline{Y}_{1,\Box}$ and cover it.
Let $T'_{1,\Box}$ be a $\mathbf{Z}/p\mathbf{Z}$-torsor over U'_1 obtained as the restriction of a $\mathbf{Z}/p\mathbf{Z}$-torsor over $\mathbf{P}^1$ ramified only above $z_{\Box,0}$ and $z'_{\Box,l}$. It is trivial over $U'_{1,\Box} \cap U'_{2,\Box}$, and, by choosing such a trivialization, one gets by descent a $\mathbf{Z}/p\mathbf{Z}$-torsor $T'_\Box$ over $\overline{Y}_{1,\Box}$, ramified only over $z_{\Box,0}$ and $z_{\Box,l}$.
$T_\Box \to \overline{Y}_{1,\Box}$ is split over x_i (with $i \in [0, mn-1]$) if and only if $|q_\Box|^{\frac{i}{n}} \in [|q_\Box|^{\frac{l}{n}} p^{\frac{p}{p-1}}, p^{-\frac{p}{p-1}}]$, *i. e.*

$$i \in I_{2,\Box} := [\frac{np}{v_p(q_\Box)(p-1)}, l - \frac{np}{v_p(q_\Box)(p-1)}].$$

There is such an integer thanks to the assumption about l, m and n because

$$\lg(I_{2,\Box}) = l - 2\frac{np}{v_p(q_\Box)(p-1)} \geqslant 1$$

.

Let finally $T''_\Box$ be the essentially unique connected topological covering of degree p of $\overline{Y}_\Box$ and let us choose an isomorphism from $\mathbf{Z}/p\mathbf{Z}$ onto its Galois group so that it becomes a $\mathbf{Z}/p\mathbf{Z}$-torsor. Let us then show that $T_\Box, T'_\Box$, and $T''_\Box$ constitute a base of $V_\Box$.
Let i be an integer in $I_{1,\Box}$. As $I_{1,\Box} \cap I_{2,\Box} = \emptyset$ and as T'' is everywhere split, if $aT_\Box + bT'_\Box + cT''_\Box$ (the linear combination is in the sense of the structure of vector space $V_\Box$) is split over x_i, $c = 0$. By the same argument with $I_{2,\Box}$, if $aT_\Box + bT'_\Box + cT''_\Box = 0$, one gets $b = 0$, but as $T_\Box$ is not trivial either, one indeed gets that $T_\Box, T'_\Box$ and $T''_\Box$ constitute a base of $V_\Box$.

Assume now ab absurdo that $|q_\alpha| > |q_\beta|$, then $I_{1,\alpha} \subset I_{1,\beta}$.
Let $T_{0,\beta} = aT_\beta + bT'_\beta + cT''_\beta$ be the image of T_α by $(\phi^{-1})^* : V_\alpha \to V_\beta$ (indeed, phi induces such a $(\phi^{-1})^*$ according to part 4.1.1).
Let $i \in I_{1,\alpha}$, T_α is split over $x_{\alpha,i}$, so, according to section 4.1.2, $T_{0,\beta}$ is split over

$x_{\beta,i}$, and thus $c = 0$. Thus, $T_{0,\beta}$ is split over every x_i if $b = 0$ or exactly over the x_i with $i \in I_{1,\beta}$ otherwise.
Yet, according to section 4.1.2, $T_{0,\beta}$ must be split exactly over the $x_{\beta,i}$ such that T_α is split over $x_{\alpha,i}$, *i. e.* $i \in I_{1,\alpha}$.
Thus, one cannot be in the case $b = 0$ and $I_{1,\beta}$ and $I_{1,\alpha}$ must contain exactly the same integers.
But this cannot be if $n \geqslant \frac{v_p(q_\beta)v_p(q_\alpha)(p-1)}{(v_p(q_\beta)-v_p(q_\alpha))p}$, because

$$\lg(I_{1,\beta}) - \lg(I_{1,\alpha}) = 2\frac{np}{v_p(q_\alpha)(p-1)} - 2\frac{np}{v_p(q_\beta)(p-1)} \geqslant 2.$$

□

Remark:
Assume $p \neq 2$. Let $\{z_{\Box,1}, \cdots, z_{\Box,4}\}$ be four rational points in $\mathbf{P}^1_{\mathbf{C}_p}$, and let ϕ be an isomorphism between the $\pi_1^{\text{temp}}(\mathbf{P}^1\backslash\{z_{\Box,1}, \cdots, z_{\Box,4}\})$, which identifies $z_{\alpha,i}$ to $z_{\beta,i}$ by the identification of the cusps of the graphs of the stable reduction (we assume that the curves have bad reduction, and that the length of the edge of the graph is $l_\Box$). Let $E_\Box$ be the unique $\mathbf{Z}/2\mathbf{Z}$ covering of $\mathbf{P}^1$ ramified over $\{z_{\Box,1}, \cdots, z_{\Box,4}\}$ and only over those points (the subgroups of index 2 corresponding to $E_\Box$ thus maps to each other). $E_\Box$ is a Tate curve with $|q_\Box| = 2l_\Box$. By theorem (4.4), $|q_\alpha| = |q_\beta|$ so $l_\alpha = l_\beta$, which proves again theorem (4.3) without assuming that the points are in $\mathbf{Q}_p^{\text{nr}}$.
If $p = 2$ and $l_\Box > 4$, then $E_\Box$ is also a Tate curve, $|q_\Box| = 2l_\Box - 8$ and the previous argument works as well.

4.4 Case of a Mumford curve

4.4.1 Reminder on Mumford curves and currents

Let X be a Mumford curve of genus g over $\overline{K}$, let $\Omega \subset \mathbf{P}^1$ be its universal topological covering, and $\Gamma = \mathrm{Gal}(\Omega/X)$, so that $X = \Omega/\Gamma$.
Let Φ be the retraction of Ω, as a Berkovich space, onto its skeleton $\mathbb{T} = \mathbb{T}(\Omega)$. For $z, z' \in \Omega$, let us set

$$d(z,z') = \sup_{x_1,x_2 \in \mathbf{P}^1\backslash\Omega} |v_p(\frac{z'-x_1}{z-x_1}\frac{z-x_2}{z'-x_2})|$$

which is invariant under homographies. Moreover $d(z,z')$ only depends on $\Phi(z)$ and $\Phi(z')$, and it is nothing else than the distance between $\Phi(z)$ and $\Phi(z')$ for the usual metric structure on the tree of Ω (that we will also denote by d, so that $d(z,z') = d(\Phi(z),\Phi(z'))$).
If $z \in \Omega$ and $\lambda > 0$, let $U_{z,\lambda}$ denote $\{z \in \Omega / d(z,z') \leqslant \alpha\}$. It is an affinoid subspace of Ω.

Let $\mathcal{L} = \mathbf{P}^1\backslash\Omega$, a compact subset of $\mathbf{P}^1$. In [8, 1.8.9], Fresnel and van der Put define a *measure* on a profinite topological space Z as a function μ : {compact open subsets of Z} $\to \mathbf{Z}$ such that $\mu(U_1 \cup U_2) + \mu(U_2 \cap U_1) = \mu(U_1) + \mu(U_2)$ for every compact open subsets U_1, U_2 of Z. The group of measures on Z such that $\mu(Z) = 0$ is then denoted by $M_0(Z)$.
One can then associate to $f \in O(\Omega)^*$ a measure μ_f on $\mathcal{L}$. Following [8], we will also call hole of an affinoid U subspace of $\mathbf{P}^1$ every connected component of the

complement of this affinoid, and will note $t(U)$ the set of its holes.
One gets the following exact sequence of groups ([8, Prop. 1.8.9]):

$$1 \to \overline{K}^* \to O(\Omega)^* \to M_0(\mathcal{L}) \to 0$$

More precisely, according to [8, 1.8.10 ex. β], if $a, b \in \mathcal{L}$ and $f : z \mapsto \frac{z-a}{z-b}$, $\mu_f = \delta_a - \delta_b$.
For a general $f_0 \in O(\Omega)^*$ and $\mu = \mu_f$, μ is a weak limit of a sequence of $\mathbf{Z}$-linear combinations $(\mu_k)_{k\in\mathbf{N}}$ of Dirac measures. Then, if $\mu_k = \sum n_i(k)\delta_{a_i(k)}$, let $f_k = \prod(1 - \frac{a_i(k)}{z})^{n_i(k)}$. Then $\mu_{f_k} = \mu_k$, and (f_k) tends uniformly on every affinoid to an invertible function g. Then $\mu_g = \mu$ and thus $g = \lambda f$ with $\lambda \in \overline{K}^*$.

According to [16, prop. 1.1], one also has the following exact sequence (where $C(\mathbb{T})$ denotes the group of currents with integral coefficients on $\mathbb{T}$):

$$1 \to \overline{K}^* \to O(\Omega)^* \to C(\mathbb{T}) \to 0$$

which gives, with the previous one an isomorphism $M_0(\mathcal{L}) \to C(\mathbb{T})$.
One can describe this isomorphism in the following way : if e is an oriented edge of $\mathbb{T}$, $\mathbf{P}^1\backslash\Phi^{-1}(e)$ has two connected components, and $\Phi^{-1}(e) \cap \mathcal{L} = \emptyset$ so one gets a partition of $\mathcal{L}$ by two open subsets $\mathcal{L}_1(e)$ at the beginning of e and $\mathcal{L}_2(e)$ at the end of e (one has for $\mu \in M_0(\mathcal{L})$, $\mu(\mathcal{L}_1(e)) = -\mu(\mathcal{L}_2(e))$. Then $\mathcal{C}(e) = \mu(\mathcal{L}_2(e))$.
more generally, $\mathbb{K}$ is a finite connected subgraph of $\mathbb{T}$ (containing at least one edge), $\Phi^{-1}(\mathbb{K})$ is an affinoid contained in Ω. There is a natural bijection between $t(\Phi^{-1}(\mathbb{K}))$ and the set of end points of $\mathbb{K}$. Then, if $\mu \in M_0(\mathcal{L})$, $C \in t(\Phi^{-1}(\mathbb{K}))$ and e is the only oriented edge of K ending at the end point of $\mathbb{K}$ corresponding to C, then $C \cap \mathcal{L} = \mathcal{L}_2(e)$ and thus $\mathcal{C}(e) = \mu(C \cap \mathcal{L})$. In particular $\mu(C \cap \mathcal{L}) = 0$ for every $C \in t(\Phi^{-1}(\mathbb{K}))$ if and only if $\mathcal{C}(e) = 0$ for every terminal edge of $\mathbb{K}$, if and only if $\mathcal{C}$ is zero on $\mathbb{K}$.

Let Θ denote the group of theta functions of X, that is the group of $f \in O(\Omega)^*$ such that for every $\gamma \in \Gamma$, $z \mapsto f(\gamma z)/f(z)$ is a constant function (this means that the corresponding current is Γ-equivariant). Then, one has the following exact sequence :

$$1 \to \overline{K}^* \to \Theta \to C(\Gamma) \to 0$$

and thus $\Theta/\overline{K}^*$ is a free $\mathbf{Z}$-module of rank g.
One deduces from [15, prop. 2.1] that :

Proposition 4.5 *for every $n \geq 2$ and every $\mathbf{Z}/n\mathbf{Z}$-torsor Y over X, there exists an element θ in Θ, unique modulo $\overline{K}^*\Theta^n$ such that $Y \times_X \Omega = \Omega[f]/(f^n = \theta)$ where $\Omega[f]/(f^n = \theta)$ denotes the pullback of the $\mathbf{Z}/n\mathbf{Z}$-torsor $\mathbf{G}_m \overset{z\mapsto z^n}{\to} \mathbf{G}_m$ along $\Omega \overset{\theta}{\to} \mathbf{G}_m$.*
Conversely, for every θ in Θ, there exists a $\mathbf{Z}/n\mathbf{Z}$-torsor Y over X such that $Y \times \Omega = \Omega[f]/(f^n = \theta)$.

Proof : Follow the notations of [15, section 2]. Then Ω_* is a connected component of $Y \times_X \Omega$.

Suppose first that $Y \times_X \Omega$ is connected, so that $\Omega_* = Y \times_X \Omega$. Then, according to [15, prop. 2.1] there is a unique lattice T in $(\Theta/\overline{K}^*) \otimes \mathbf{Q}$ containing $\Theta/\overline{K}^*$ such that, as a X-covering, $\Omega_* = \Omega(T) := \Omega \times_{\operatorname{Spec} \overline{K}[\Theta/\overline{K}^*]} \operatorname{Spec} \overline{K}[T]$. Then $T/(\Theta/\overline{K}^*)$ is isomorphic to $\mathbf{Z}/n\mathbf{Z}$, and choosing a $\bar{f}$ generator of $T/(\Theta/\overline{K}^*))$ amounts to choosing a $\mathbf{Z}/n\mathbf{Z}$-torsor structure on Ω_*.
If one takes $\bar{f}^i$ with $i \in (\mathbf{Z}/n\mathbf{Z})^*$ as another generator, the torsor one gets is $\Omega_*^{\otimes i}$. Thus by changing the generator, one can get all the $\phi(n)$ $\mathbf{Z}/n\mathbf{Z}$-torsor structure on the covering Ω_* of Ω.
If $\bar{f}$ is the generator corresponding to the $\mathbf{Z}/n\mathbf{Z}$-torsor structure on $Y \times_X \Omega$, then θ is any lifting of $\bar{f}^n \in T^n/(\Theta/\overline{K}^*)^n$.
In the general case Ω_* acquires the structure of a $\mathbf{Z}/m\mathbf{Z}$-torsor over Ω with $m|n$, and as before one can find a unique θ_0 modulo $\overline{K}^*\Theta^m$ such that $\Omega_* = \Omega[f]/(f^m = \theta_0)$. Then $Y \times_X \Omega = \operatorname{Ind}_{\mathbf{Z}/m\mathbf{Z}}^{\mathbf{Z}/n\mathbf{Z}} \Omega_*$, and thus $Y \times_X \Omega = \Omega[f]/(f^n = \theta_0^{m/n})$, and $\theta = \theta_0^{m/n}$.
The second statement comes from the fact that, if Ω_* is a connected component of $\Omega[f]/(f^n = \theta)$, $\operatorname{Gal}(\Omega_*/X)$ is (non canonically) isomorphic to the direct product of $\operatorname{Gal}(\Omega_*/\Omega)$ and Γ (according to [15, section 2, intro.]). Thus Ω_* can descend (non canonically) to X by considering $Y_0 = \Omega_*/N$ where N is some complement of $\operatorname{Gal}(\Omega_*/\Omega)$ and $\operatorname{Gal}(\Omega_*/X)$ (and thus $\Omega[f]/(f^n = \theta)$ by taking a direct sum of Y_0s. □

Remark: One could also show it by considering $\widetilde{J} = \mathcal{H}om(\Theta/\overline{K}^*, \mathbf{G}_m) \to J$ where J is the Jacobian variety of X and $\widetilde{J}$ is its universal topological covering, and by showing that $\pi_1^{\mathrm{alg}}(\widetilde{J})$ is a direct summand of $\pi_1^{\mathrm{alg}}(J)$.

4.4.2 Preliminary results on the ramifications of torsors corresponding to currents

Proposition 4.6 *Let $z \in \Omega, \lambda > 0$. Let $f \in O(\Omega)^*$ such that $f(z) = 1$. Let μ be the measure on $\mathcal{L}$ corresponding to f and let us assume $\mu(C \cap \mathcal{L}) = 0$ for every hole of $U_{z,\lambda}$.
Then $\forall z' \in U_{z,\lambda}$, $|f(z') - 1| \leqslant p^{d(z,z')-\lambda}$*

Proof : To simplify, assume $z = \infty$.
According to [8, 1.8.10 ex. β] and [8, prop. 1.8.9.(i)], $f = \lim f_k$ uniformly on every affinoid of Ω (in particular over $U_{z,\lambda}$) where f_k is of the following form

$$f_k(z') = \prod_{i=1}^{s_k} (1 - \frac{x_{i,k}}{z'})^{n_{i,k}},$$

and $\mu_k = \sum_i n_{i,k}\delta_{a_i(k)} \to \mu$.
For k big enough, $\mu_k(C) = 0$ for every hole C of $U_{z,\lambda}$ and $|f - f_k|_{U_{z,\lambda}} \leqslant p - \lambda$.
We thus only have to prove the result for f_k, which is a product of function such as $g : z' \mapsto z' = (1 - \frac{x_1}{z'})(1 - \frac{x_2}{z'})^{-1} = \frac{z'-x_1}{z'-x_2}$ with x_1 and x_2 in the same hole C of $U_{z,\alpha}$.
We thus only have to prove the result for g, which is easily seen. □

Corollary 4.7 *Let $f \in O(\Omega)^*$ such that $f(z) = 1$. Let U an affinoid of Ω such that for every hole C of U, $\mu(C) = 0$.
Assume that, for all $z'' \in \Phi^{-1}([\Phi(z), \Phi(z')]), U_{z'',\lambda} \subset U$.
Then $|f(z') - 1| \leqslant p^{-\lambda}$.*

Proof : Let $\epsilon > 0$. Let $(z_i)_{i=0\cdots n}$ such that $z_0 = z$, $z_n = z'$, $\Phi(z_i) \in [\Phi(z), \Phi(z')]$ and $d(\Phi(z_i), \Phi(z_{i+1})) \leqslant \epsilon$.
Then, according to the previous proposition $|\frac{f(z_{i+1})}{f(z_i)} - 1| \leqslant p^{\epsilon-\lambda}$.
Thus $|f(z'') - 1| \leqslant \sup |f(z_{i+1}) - f(z_i)| \leqslant p^{\epsilon-\alpha}$. Ones get the result by letting ϵ going to 0. □

Corollary 4.8 *Let f as previously, U such that $\mu(C) = 0$ for every hole C of U. Let e a positive integer. Let $\lambda > e + \frac{1}{p-1}$, let $Y \to \Omega$ be finite covering obtained by pulling back $\mathbf{G}_m \overset{z\mapsto z^{p^e}}{\to} \mathbf{G}_m$ along $f : \Omega \to \mathbf{G}_m$. Let $V \subset U$ be such that $\forall z \in V, U_{z,\lambda} \subset U$.*
Then Y is split over V.

Proof : We may assume V connected, because we only have to prove the result for every connected component of V. Let $z \in V$. Multiplying f by a constant, which do not change Y, we may assume that $f(z) = 1$.
From the previous corollary, $f(V) \subset D(1, p^{-\lambda})$. But, according to lemma (4.2), $\mathbf{G}_m \overset{z\mapsto z^{p^e}}{\to} \mathbf{G}_m$ is split over $D(1, p^{-\lambda})$, which ends the proof. □

Proposition 4.9 *Let $\mathcal{C}$ a current on $\mathbb{T}(\Omega)$, corresponding to an invertible function f on Ω. Let a a vertex of $\mathbb{T}$ such that the restriction $\mathcal{C}_a$ of $\mathcal{C}$ the star of a is not zero modulo n. Then, if $Y \to \Omega$ is the finite covering obtained by pullback of $\mathbf{G}_m \overset{z\mapsto z^n}{\to} \mathbf{G}_m$ along $f : \Omega \to \mathbf{G}_m$, $Y \to \Omega$ is not split over a.*

Proof : $Y \to \Omega$ is split over a if and only if there exists $f_1 \in \mathcal{O}_{\Omega,a}$ such that $f_1^n = f_{|\mathcal{O}_{\Omega,a}}$.
By multiplying f by a constant, we may assume $|f|_a = 1$.
If f_1 exists, by looking at the residue field $\tilde{k}_{\Omega,a} \simeq \tilde{k}(X)$, $\tilde{f}_1^n = \tilde{f}$. If $\tilde{f}_1(z) = \lambda \prod (z - a_i)$, $\tilde{f} = \lambda^n \prod (z - a_i)^n$, and so all the poles and zeros are of order a muliple of n, which ends the proof. □

4.4.3 The metric graph of the reduction and the tempered fundamental group

Assume now that X_α and X_β soient are two Mumford curves over $\overline{K}$, but are pullback of curves over K (so that we may use [13, ex. 3.10]), and that there is an isomorphism $\phi : \pi_1^{\text{temp}}(X_\alpha) = \pi_1^{\text{temp}}(X_\beta)$, which thus inducts an isomorphism of graphs $\mathbb{G}(X_\alpha) \to \mathbb{G}(X_\beta)$. Hence an isomorphism $\mathbb{T}(\Omega_\alpha) \simeq T(\Omega_\beta)$.

Theorem 4.10 *The isomorphism $\mathbb{G}_\alpha \to \mathbb{G}_\beta$ of graph induced by an isomorphism $\pi_1^{\text{temp}}(X_\alpha) \to \pi_1^{\text{temp}}(X_\beta)$ of topological groups is in fact an isomorphism of metric graphs.*

Remark: Suppose X_α and X_β are projective lines minus four points $a_\Box, b_\Box, c_\Box, d_\Box$ with $v_p(a_\Box, b_\Box, c_\Box, d_\Box) = e_\Box$with an isomorphism ϕ between their tempered fundamental groups, and suppose $p \neq 3$. One can consider a covering X'_α of order three of X_α such that the restriction to each irreducible component of the stable reduction is connected but it is split over the double point of the stable reduction (and let X'_β be the covering of X_β corresponding to X'_α by ϕ, it satisfies the

same properties). Then $\overline{X}'_\square$ is a Mumford curve (the corresponding covering of each irreducible component of the stable reduction of $X_\square$ is ramified in only two points, so it is a covering by a projective line) whose tree has three edges, each of length $e_\square$. Thus, by what has just been proven $e_\alpha = e_\beta$, which ends the proof for punctured lines.
One could in fact do quite the same for more general punctured Mumford curve, by considering, for an edge e of the graph, a tamely ramified covering by a Mumford curve such that there is an edge e' over e which is actually an edge of the graph of the stable reduction of the compactification of this covering.

Proof : A *loop* of a graph is a cyclic sequence of oriented edges of the graph such that the end of an edge is the beginning of the following edge, and which never goes by the same vertex or (unoriented) edge twice.
Let C be a loop of this graph, and note $\lg_\square(C)$ the length of C with respect to the metric $d_\square$ on $\mathbb{G}$. Let $\widetilde{C}$ be the universal covering of C, let $\widetilde{C} \to \mathbb{T}$ a lifting of $\widetilde{C} \to \mathbb{G}$ and let z_0 a vertex of $\mathbb{T}$ which belongs to $\widetilde{C}$, let us number then $(z_i)_{i\in\mathbf{Z}}$ the vertices of $\widetilde{C}$ with the same image that z_0 in $\mathbb{G}$. Let L be another loop (we choose an orientation on it) of $\mathbb{G}$ (there must be another loop since $g > 1$), let $\widetilde{L}$ be a lifting of L to $\mathbb{T}$, let $r_\square = d_\square(\widetilde{L}, z_0)$ (we may assume, by changing the numbering of the z_i that for $n \geqslant 0$, $d_\square(\widetilde{L}, z_n) = r_\square + n \lg_\square(C)$) and let z'_0 the point of $\widetilde{L}$ the nearest from z_0 (this does not depend on $\square$).

Let us note, for z a vertex of $\mathbb{T}$, F_z the connected component of $\Omega\backslash\{\text{open edges of } \widetilde{L}\}$ which contains z.

Let $e \geqslant 1$ be an integer.
Let K_0 be a connected finite subgraph of $\mathbb{T}$ containing z_1 (so that $\Phi^{-1}(|K_0|)$ is compact by properness of Φ).
Let $K_\square$ a connected finite subgraph (coinciding for $\square = \alpha$ and $\square = \beta$) $\mathbb{T}(\Omega_\square)$ such that for every $z \in \Phi^{-1}(|K_0|)$ $U_{z,e+2} \subset \Phi^{-1}(|K|)$ and such that, for every vertex z of $\widetilde{L} \cap K_\square$, $\{z' \in F_z | d(z', z) \leqslant e + 2\} \subset \Phi^{-1}(|K_\square|)$ (in particular, if z' is a vertex of the boundary of K in $\mathbb{T}$ which is not one of the end points of the segment $\widetilde{L} \cap K$, $d(z', \widetilde{L}) \geqslant e + 2$). Let $K'_\square$ be a compact subgraph of $\mathbb{T}$ coinciding for $\square = \alpha$ and $\square = \beta$ such that $\Phi_\square^{-1}(|K'_\square|)$ contains $U_{z,e+2}$ for every z in $\Phi^{-1}(K_\square)$.

Let $\Gamma = \mathrm{Gal}(\mathbb{T}/\mathbb{G})$, let $H = \mathrm{Stab}(\widetilde{L}) (\simeq \mathbf{Z})$ and let Γ' be a subgroup of finite index of Γ such that, for every $g \neq 1 \in \Gamma'$, $\Phi_\square^{-1}(|K'|) \cap g \cdot \Phi_\square^{-1}(|K'|) = \emptyset$ and for every $g \in \Gamma\backslash H$, $d_\square(g \cdot \widetilde{L}, \widetilde{L}) > \mathrm{diam}_\square(|K'|)$($A = \{g \neq 1 \in \Gamma | \Phi^{-1}(|K'|) \cap g \cdot \Phi^{-1}(|K'|) \neq \emptyset\}$ is finite by compactness of $\Phi^{-1}(|K'|)$. So, as Γ is residually finite, there exists Γ'_1 of finite index in Γ which does not intersect A. $B = \{g \in \Gamma/H - H | d_\square(g \cdot \widetilde{L}, \widetilde{L}) \leqslant \mathrm{diam}_\square(|K'|)\}$ is also finite, and as $\overline{H} \cap \Gamma = H$ in the profinite completion of Γ, there exists Γ'_2 of finite index in Γ containing H such that $\Gamma'_2 \cap B \cdot H = \emptyset$. We may then chose $\Gamma' = \Gamma'_1 \cap \Gamma'_2$). Let $H' = H \cap \Gamma'$. Let $X'_\square = \Omega_\square/\Gamma'$: its a finite topological covering of $X_\square$, and the isomorphism $\phi : \pi_1^{\mathrm{temp}}(X_\alpha) \simeq \pi_1^{\mathrm{temp}}(X_\beta)$ inducts an isomorphism $\phi' : \pi_1^{\mathrm{temp}}(X'_\alpha) \simeq \pi_1^{\mathrm{temp}}(X'_\beta)$.

Let $\mathcal{C}_0$ be the current on $\mathbb{T}$ with $\mathcal{C}_0(e) = +1$ if e is an edge of $\widetilde{L}$ (and e has the good orientation) and 0 otherwise (except if e is an edge of $\widetilde{L}$ with the wrong orientation, in which case $\mathcal{C}_0(e) = -1$) : this current is invariant under H. Let $\mathcal{C}_\alpha = \sum_{g \in \Gamma'/H'} g \cdot \mathcal{C}_0$. It is a current on $\mathbb{T}$, invariant under Γ' and which coincides with $\mathcal{C}_0$ on K'.
Let $f_\alpha \in O(\Omega_\alpha)^*$ be the corresponding invertible function with $f_\alpha(z_{0,\alpha}) = 1$, and let X''_α be a $(\mathbf{Z}/p^e\mathbf{Z})$-torsor of X'_α corresponding to that current, that is to say such that its pullback $\widetilde{X}''_\alpha$ to Ω_α is isomorphic to $\Omega_\alpha \times_{\mathbf{G}_m} \mathbf{G}_m \to \Omega_\alpha$ where the fiber product is taken, on the left side, along f_α and on the right side along $z \mapsto z^{p^e}$. Let $X''_\beta = \phi'^* X''_\alpha$ (do not forget that X''_β has no reason to correspond to the current $\mathcal{C}_\alpha$).
Let also $f_{0,\square} \in O(\Omega_\square)^*$ be the invertible function corresponding to the current $\mathcal{C}_0$ and let $\widetilde{X}_{0,\square}$ be the corresponding $(\mathbf{Z}/p^e\mathbf{Z})$-torsor on $\Omega_\square$. Recall that this torsor is split over a point $z \in \mathbb{T}(\Omega_\square)$ of $\Omega_\square$ if and only if $d_\square(z, \widetilde{L}) > e + \frac{1}{p-1}$.

According to corollary (4.8) applied to $U = \Phi^{-1}(|K'|)$ and to $V = \Phi^{-1}(|K|)$, the torsor $\widetilde{X}''_\alpha - \widetilde{X}_{0,\alpha}$ on Ω_α, which corresponds to the current $\mathcal{C}_\alpha - \mathcal{C}_0$ which is zero over K', is split over $\Phi^{-1}(|K|)$ since for every $z \in \Phi^{-1}(|K|)$, $U_{z,e+2} \in \Phi^{-1}(|K'|)$. Thus, for $z \in K$, $\widetilde{X}''_\alpha$ is split if and only if $\widetilde{X}_{0,\alpha}$ is, if and only if $d_\alpha(z, \widetilde{L}) > e + \frac{1}{p-1}$.
In particular $\widetilde{X}''_\alpha$ is split over the vertices of the boundary of K which are not the end points of $K \cap \widetilde{L}$. Thus, according to the result of section 4.1.2 applied to X''_α and X''_β (as $\Omega_\square \to X'_\square$ is a topological covering, $\widetilde{X}''_\square \to \Omega_\square$ is split over a point if and only if $X''_\square \to X'_\square$ over the image of that point), $\widetilde{X}''_{0,\beta}$ is also split over the vertices of the boundary of K which are not the end points of $K \cap \widetilde{L}$.

Let $\mathcal{C}_\beta$ be a current on $\mathbb{T}(\Omega_\beta)$ corresponding to the $\mathbf{Z}/p^e\mathbf{Z}$-torsor $\widetilde{X}''_\beta$ (the current corresponding to $\widetilde{X}''_\beta$ is well defined only modulo p^e). According to proposition (4.9), 1 the restriction of $\mathcal{C}_\beta$ to the star of a vertex of K which is not an end point of $K \cap \widetilde{L}$ is zero modulo p^e. One deduces from it that, modulo p^e, the restriction to K of $\mathcal{C}_\beta$ must be congruent to the restriction of $a\mathcal{C}_0$. By adding to $\mathcal{C}_\beta$ a current which is a multiple of p^e, we may assume that $\mathcal{C}_\beta - a\mathcal{C}_0$ is zero on K (because every current with boundary on K, that is to say that respect Kirchhoff's law in every vertex of the interior of K but with no condition on the boundary of K, can be extended in a current on the whole $\mathbb{T}$).
Thus, by applying corollary (4.8) to $U = \Phi^{-1}(|K|)$ and to $V = \Phi^{-1}(K_0)$, one may deduce that $\widetilde{X}''_\beta - a\widetilde{X}_{0,\beta}$ is split over $|K_0|$, so if z is a vertex of K_0, $\widetilde{X}''_\beta$ is split over z if and only if $\widetilde{X}_{0,\beta}$ is (a is necessarily non zero modulo p^e because $\widetilde{X}''_\beta$ cannot be split over z'_0), if and only if $d_\beta(z, \widetilde{L}) > e + \frac{1}{p-1}$, according to lemma (4.2).

Therefore, $d_\beta(z, \widetilde{L}) > e + \frac{1}{p-1}$ if and only if $d_\beta(z, \widetilde{L}) > e + \frac{1}{p-1}$ if and only if $d_\alpha(z, \widetilde{L}) > e + \frac{1}{p-1}$ for every vertex $z \in K_0$.
As one may choose K_0 as big as one wants, one may deduce that for every z of $\mathbb{T}$, $d_\alpha(z, \widetilde{L}) > e + \frac{1}{p-1}$ if and only if $d_\beta(z, \widetilde{L}) > e + \frac{1}{p-1}$, and this for every integer $e \geqslant 1$.
Thus $\max(1, \lceil d_\beta(z, \widetilde{L}) - \frac{1}{p-1} \rceil) = \max(1, \lceil d_\alpha(z, \widetilde{L}) - \frac{1}{p-1} \rceil)$.

By applying it to $(z_i)_{i\geqslant 0}$, one gets that for every $i \geqslant 0$,

$$\max(1, \lceil i \lg_\alpha + r_\alpha - \frac{1}{p-1} \rceil) = \max(1, \lceil i \lg_\alpha + r_\alpha - \frac{1}{p-1} \rceil).$$

One may deduce from this that for every loop C of $\mathbb{G}$ (and of every topological covering of $\mathbb{G}$),

$$\lg_\alpha(C) = \lg_\beta(C).$$

One may conclude with the help of proposition (A.1). $\square$

A A combinatorial result

To end the proof of theorem (4.10), we will need to prove that one can recover from the length of all the loops of every finite covering of $\mathbb{G}$ the length of every edge of $\mathbb{G}$. We thus have to prove a general result on graphs such that the valency of every edge is at least 3.
To prove this, we will work by induction on the number of edges of the graph. However, by removing an edge of our graph, one does not in general get a graph such that the valency of every edge is at least three, and we may have to concatenate two edges or withdrawing an other edge to apply our induction assumption (but there will only be a few edges whose length cannot be recovered directly by the induction assumption). To exhibit enough loops that go through some edges we do not know yet the length thanks to the induction assumption, we will have to distinguish many cases depending on the number of connected components of the subgraph of all the edges whose length is already known.

Proposition A.1 *Let $\mathbb{G}$ be finite graph such that the valency of every vertex is at least 3. Let $f : \{\text{edges of } \mathbb{G}\} \to \mathbf{R}$ be any function. Let us denote also by f the induced function on the set of edge of a (topological) covering of $\mathbb{G}$. Let us set, for C a loop of a covering of $\mathbb{G}$, $f(C) = \sum_{x\in\{\text{edges of } C\}} f(x)$.*
If $f(C) = 0$ for every loop C of every covering of $\mathbb{G}$, then $f = 0$.

Proof : Remark that if $\mathbb{G}$ is a finite graph such that the valency of every vertex is at least 3 and if $\mathbb{H}$ is a connected subgraph such that the number of half-edges of $\mathbb{G}\backslash\mathbb{H}$ which ends in $\mathbb{H}$ is less than 3, then $\mathbb{H}$ is not a tree (if $\mathbb{H}$ is a tree with at least one edge, $\mathbb{H}$ has at least two vertices of valency 1, and thus one already has 4 half-edges of $\mathbb{G}\backslash\mathbb{H}$ which must end in one of those two vertices; if $\mathbb{H}$ is only a vertex, it is equally obvious).

Let us go back to our business. We will proceed by induction on the number of edges of $\mathbb{G}$.
Thus let $(\mathbb{G}, f)$ be a graph with $n \geqslant 1$ edges and a function f on the set of edges of $\mathbb{G}$ which verify the hypotheses of the proposition, and assume the proposition to be true if $\mathbb{G}$ has less than n edges.
We may assume $\mathbb{G}$ to be connected (otherwise we may use the induction hypothesis to the various connected components to conclude).
Let e be an edge of $\mathbb{G}$, and start with assuming that:

1. the two end points of e are different. Let (m, n) be the arities of the end points of e in $\mathbb{G}\backslash\{e\}$. By our assumption about $\mathbb{G}$, $m \geqslant 2$ and $n \geqslant 2$.

(a) If $m \geqslant 3$ and $n \geqslant 3$, $\mathbb{G}' = \mathbb{G} \backslash \{e\}$ is still a graph such that the valency of every vertex is at least 3, thus one may apply our induction hypothesis to $\mathbb{G}'$ and to $f: f(x) = 0$, for every edge $\mathbb{G}$ other than e.

   i. If $\mathbb{G}'$ is connected, then one may find a loop C of $\mathbb{G}$ that goes through e, and then $f(e) = f(C) = 0$, which implies the result.

   ii. If $\mathbb{G}'$ has two connected components A and B (they cannot be trees according to the remark at the beginning of the proof), one may consider two coverings of order 2 A' and B' of A and B respectively, that one patches in a covering $\mathbb{G}'$ of G of order 2. Then there exist a loop C of $\mathbb{G}'$ passing through the two edges over e.
   Then $2f(e) = f(C) = 0$, which gives the wanted result.

(b) If $m \geqslant 3$ and $n = 2$ (or the other way), then let a and b be the two edges starting at the second end point of e (if a and b are in fact the two half-edges of a single edge, the graph has then the same structure as the one of case 2.(c).i, where it will be treated; thus we will assume here that a and b are two different edges). Let $\mathbb{G}'$ the graph obtained from $\mathbb{G}$ by withdrawing e and concatenating a and b in a single edge that we will note $a + b$ (and we will define $f(a+b) = f(a)+f(b)$). Then $(\mathbb{G}', f)$ satisfies the required conditions and so $f(x) = 0$ for every edge of $\mathbb{G}$ other than a, b and e. Depending on the number of connected components of $\mathbb{G}'' = \mathbb{G} \backslash \{a, b, e\}$, we will distinguish several cases:

   i. If $\mathbb{G}''$ has only one connected component, then one may contract $\mathbb{G}''$ to a point to get a graph $\mathbb{G}_1$ with three edges (indeed, every loop C_1 of $\mathbb{G}_1$ may be lifted to a loop C of $\mathbb{G}$ as $\mathbb{G}''$ is connected, and $f(C_1) = f(C)$ since $f = 0$ over $\mathbb{G}''$), and thus $f(a)+f(b) = 0$, $f(a) + f(e) = 0$ and $f(b) + f(e) = 0$, which gives the wanted result.



   ii. If $\mathbb{G}''$ has two connected components A and B as in the picture (now a, b and e will play the same role and the proof will be the same if they are exchanged), then we start by considering a connected covering $\mathbb{G}_1$ of order 2 of $\mathbb{G}$ such that its restrictions A' and B' to A and to B are connected (there exist such a covering as A and B cannot be trees according to the remark at the beginning of the proof), then one may contract A' and B' into a graph G_2. One gets $f(b) + f(e) = 0$, $2f(a) + 2f(b) = 0$, $2f(a) + 2f(e) = 0$, which implies what we wanted.

   iii. If $\mathbb{G}''$ has three connected components A, B and C, then we start by considering a connected covering $\mathbb{G}_1$ of order 2 of $\mathbb{G}$ such that







its restrictions A', B' and C' to A, B and C are connected, then one may contract A', B' and C'. One may deduce that $2f(a) + 2f(b) = 0, 2f(b) + 2f(e) = 0, 2f(e) + 2f(a) = 0$, which implies the result.







(c) If $m = n = 2$, then let a, b be the two (half-)edges ending at one end point of e and c, d the (half-)edges ending at the other end point of e. Let $\mathbb{G}'$ be the graph obtained from $\mathbb{G}$ by withdrawing e and concatenating a and b into $a + b$, and c and d into $c + d$. If one defines $f(a + b) = f(a) + f(b)$ and $f(c + d) = f(c) + f(d)$, (G', f) satisfies assumptions of the proposition, and so, by the induction hypothesis, $f(x) = 0$ for every edge x of $\mathbb{G}$ else than a, b, c, d and e. Let $\mathbb{G}'' = \mathbb{G} \backslash \{a, b, c, d, e\}$. Depending on the connected components of $\mathbb{G}''$, we will distinguish several cases :

i. If $\mathbb{G}$ has two connected components, A containing one end point of a and of b, and B containing one end point of c and of d (if (a, b) or (c, d) are the two half-edges of a single edge, the graph has the same structure as in case 2.(c).ii.B; if (a, b) and (c, d) both make single edges, then the structure is the one of the degenerate case of 2.(c).(ii)), then start by considering a connected covering $\mathbb{G}_1$ of order 2 of $\mathbb{G}$ such that the restrictions A' and B' to A and B are connected, and then contract A' and B'. One gets, for example, that $f(c)+f(d), f(a)+f(b), 2(f(a)+f(e)+f(d)), 2(f(a)+f(e)+f(c), 2(f(b) + f(e) + f(d)) = 0$, which implies the result.





ii. If $\mathbb{G}$ has two connected components, A containing an end point of a and c, and B containing an end point of b and d, then consider as usual a connected covering $\mathbb{G}_1$ of order 2 of $\mathbb{G}$ such that the restrictions A' and B' to A and B are connected, then contract A' and B'. One gets for example that $f(a)+f(b)+f(c)+f(d) = 2(f(a)+f(e)+f(d)) = 2(f(b)+f(e)+f(c)) = 2(f(a)+f(c)) = f(b)+f(c)+f(e) = 0$, which implies the result.





If (a,c) (or symmetrically (b,d)) are the two half-edges of a single edge a, consider a connected covering $\mathbb{G}_1$ of order 2 of $\mathbb{G}$ such that the restrictions B' to B and $(a \cup e)'$ and $a \cup e$ are connected. One gets that $f(a)+f(b)+f(c) = f(e)+f(b)+f(c) = 2(f(a)+f(e)) = f(a)+f(e)+2f(b) = 0$, which implies the result.





If (a,c) and (b,c) are degenerated into two edges, then $\mathbb{G}$ only has two vertices and three arrows joining those two vertices, and the result is obvious.

iii. If $\mathbb{G}$ has two connected components, A containing an end point of a,b and of c, and B containing an end point of d. We already know by induction hypothesis applied to $\mathbb{G}'$ that $f(c)+f(d) = 0$. Now, by contracting A, one gets that $f(a)+f(b) = f(b)+f(c)+f(e) = f(a)+f(c)+f(e) = 0$, which implies the result.



iv. If $\mathbb{G}$ has three connected components, A containing an end point of a, B containing an end point of b, et C containing an end point of c and of d. Consider as usual a connected covering $\mathbb{G}_1$ of order 2 of $\mathbb{G}$ such that its restrictions A', B' and C' to A, B and C are connected, then contract A', B' and C'. One gets for example that $f(c)+f(d) = 2(f(a)+f(b)) = 2(f(a)+f(e)+f(c)) =$

$2(f(b)+f(e)+f(c)) = 2f(a)+2f(e)+f(c)+f(d) = 0$, which implies the result.



If (c,d) is degenerated in a single edge c, the graph has the same structure as in case 2.(c).ii.A.

v. If $\mathbb{G}$ has three connected components, A containing an end point of a, B containing an end point of c, and C containing an end point of b and d, then consider a connected covering $\mathbb{G}_1$ of order 2 of $\mathbb{G}$ such that its restrictions A', B' and C' to A, B and C are connected then contract A', B' and C'. One gets for example that $f(b)+f(d)+f(e) = f(b)+f(d)+f(e)+2f(a) = 2(f(a)+f(b)) = 2(f(d)+f(c)) = 2(f(c)+f(e)+f(a)) = 0$.



If (b,d) is degenerated into a single edge b, consider a covering G_1 of order 2 of G such that its restrictions to A, B and $b \cap e$ are connected, then contract A' and B'. One gets that $2f(a) + f(e) + f(b) = 2(f(b)+f(e)) = 2f(c)+f(b)+f(e) = 2(f(a)+f(b)+f(c)) = 0$.



vi. If $\mathbb{G}$ has four connected components, A containing an end point of a, B containing an end point of c, C containing an end point of b and D containing an end point of d, then consider a connected covering $\mathbb{G}_1$ of order 2 of $\mathbb{G}$ such that its restrictions A', B', C' and D' to A, B, C and D are connected then contract A', B', C' and D'. One gets for example that $2(f(b)+f(d)+f(e)) = f(b)+f(d)+f(e)+2f(a) = f(b)+f(d)+f(e)+2f(c) = 2(f(b)+f(a)) = 2(f(c)+f(d)) = 0$.



vii. If $\mathbb{G}$ has a single connected component A, then contract it. One gets that $f(a) + f(b) = f(c) + f(d) = f(a) + f(c) + f(e) = f(b) + f(c) + f(e) = f(a) + f(d) + f(e) = 0$, which implies the result.



2. the two end points of e are a same vertex, and let m be the valency of this vertex in $\mathbb{G}\backslash\{e\}$. One has $m \geqslant 1$.
   (a) Assume $m \geqslant 3$. Then $\mathbb{G}\backslash\{e\}$ satisfies the assumptions of the proposition and thus by induction hypothesis $f(x) = 0$ for every edge of $\mathbb{G}$ else than e, and as e is already a loop, $f(e) = 0$ too.
   (b) Assume $m = 2$, and let a and b be the two edges ending at the end point of e (if a and b are in fact only the two half-edges of a single edge, $\mathbb{G}$ is only the wedge of two loops, and the result is obvious). Let $\mathbb{G}'$ be the graph obtained from $\mathbb{G}$ by withdrawing e and by concatenating a and b in an edge $a+b$ and define $f(a+b) = f(a) + f(b)$. Then $\mathbb{G}'$ satisfies the assumptions of the proposition and thus, by induction hypothesis, $f(x) = 0$ for every edge x else than e, a et b. Depending on the connected components of $\mathbb{G}'' = \mathbb{G}\backslash\{a, b, e\}$, we will distinguish the following cases :
      i. If $\mathbb{G}''$ has a single connected component A, then consider a covering $\mathbb{G}_1$ of order 2 of $\mathbb{G}$ such that its restrictions to A and e are connected, then contract A': one gets a graph $\mathbb{G}_2$. One gets $2f(a) + f(e) = 2f(b) + f(e) = 2f(e) = 0$, which implies the result.



      ii. If $\mathbb{G}''$ has two connected components, A containing the end point of a, and B containing the end point of b. Consider a covering $\mathbb{G}_1$ of order 2 of $\mathbb{G}$ such that its restrictions A' and B' to A and B are connected and such that its restriction to e is connected too, then contract A' and B' to get a graph $\mathbb{G}_2$. One gets that

$2f(a) + f(e) = 2f(b) + f(e) = 2f(e) = 0$, which implies the result.



(c) Assume $m = 1$, and let a be the edge ending at the end point of e. Let n be the valency of the other end point of a. One must have $n \geqslant 2$.

i. Assume $n \geqslant 3$. Then $\mathbb{G}' = \mathbb{G}\backslash\{a, e\}$ satisfies the assumptions of the proposition, so, by induction hypothesis, $f = 0$ over $\mathbb{G}'$. Consider now a covering $\mathbb{G}_1$ of order 2 of $\mathbb{G}$ such that its restrictions to $\mathbb{G}'$ and to e are connected, and contract the preimage of $\mathbb{G}'$. One gets $2f(e) = 2f(a) + f(e) = 0$, which implies the result



ii. Assume $n \geqslant 2$, and let c and d be those two edges (if they only are the two half-edges of a single edge, $\mathbb{G}$ is made of two loops joined by an edge; one shows the result for this particular graph by by considering the covering of order 2 whose restrictions to the loops are connected). Let $\mathbb{G}''$ be the graph obtained from $\mathbb{G}'$ by concatenating c and d into $c + d$ and let us define $f(c+d) = f(c)+f(d)$. $\mathbb{G}''$ satisfies the assumptions of the proposition, and so $f(x) = 0$ for every edge x of $\mathbb{G}$ else than a, e, c and d (and, in fact, $f(e) = 0$ too). Let $\mathbb{G}''' = \mathbb{G}\backslash\{a, c, d, e\}$, and distinguish the following cases depending on the number of connected components of $\mathbb{G}'''$.

A. If $\mathbb{G}'''$ has two connected components, C containing the end point of c and D containing the end point of d, consider a covering $\mathbb{G}_1$ of $\mathbb{G}$ such that its restrictions C', D' and e' to C, D and e are connected, then contract C', D' and e', one gets $2f(a) + 2f(c) = 2f(a) + 2f(d) = 2f(c) + 2f(d) = 0$, which implies the result.



B. If $\mathbb{G}'''$ has a single connected component A, consider a covering $\mathbb{G}_1$ of $\mathbb{G}$ such that its restrictions A' and e' to A and e are connected, then contract A' and e', one gets $2f(a)+2f(c) = 2f(a)+2f(d) = 2f(c)+2f(d) = 0$, which implies the result.



□

References

[1] Yves André, *On a geometric description of* Gal$(\overline{\mathbf{Q}}_p/\mathbf{Q}_p)$ *and a p-adic avatar of* $\widehat{GT}$, Duke Mathematical Journal **119** (2003), 1–39.

[2] ______, *Period mappings and differential equations : From* $\mathbf{C}$ *to* $\mathbf{C}_p$, MSJ Memoirs, vol. 12, Mathematical Society of Japan, Tokyo, 2003.

[3] Vladimir G. Berkovich, *Smooth p-adic analytic spaces are locally contractible*, Inventiones Mathematicae **137** (1999), 1–84.

[4] ______, *Smooth p-adic analytic spaces are locally contractible. II*, Geometric aspects of Dwork theory, Walter de Gruyter & co, 2004, pp. 293–370.

[5] Aise Johan de Jong, *étale fundametal group of non archimedean analytic spaces*, Compositio mathematica **97** (1995), 89–118.

[6] ______, *Smoothness, semi-stability and alterations*, Publications mathématiques de l'IHES **83** (1996), 51–93.

[7] ______, *Families of curves and alterations*, Annales de l'institut Fourier **47** (1997), 599–621.

[8] Jean Fresnel and Marius van der Put, *Géométrie analytique rigide et applications*, Progress in Mathematics, vol. 18, Birkhäuser, 1981.

[9] ______, *uniformisation des variétés abéliennes*, annales de la faculté des sciences de Toulouse 5e série **S10** (1989), 7–42.

[10] Alexander Grothendieck (ed.), *Revêtements étales et groupe fondamental (SGA1)*, Lecture Notes in Mathematics, vol. 224, Berlin, Springer, 1971.

[11] J. S. Milne, *Jacobian varieties*, Arithmetic Geometry (Proc. Conference on Arithmetic Geometry, Storrs, August 1984), Springer, 1986, pp. 167–212.

[12] Shinichi Mochizuki, *The absolute anabelian geometry of hyperbolic curves*, Galois Theory and Modular Forms, Kluwer Academic Publishers, 2003, pp. 77–122.

[13] ———, *Semi-graphs of anabelioids*, Publications of the Research Institute of Mathematical Sciences **42** (2006), no. 1, 221–322.

[14] Amaury Thuillier, *Théorie du potentiel sur les courbes en géométrie analytique non archimédienne. application à la théorie d'arakelov*, Ph.D. thesis, Institut de Recherche Mathématique de Rennes, Rennes, 2005.

[15] Marius van der Put, *Etale coverings of a mumford curve*, Annales de l'Institut Fourier **33** (1983), no. 1, 29–52.

[16] Marius van der Put, *Les fonctions thêta d'une courbe de Mumford.*, Groupe de travail d'Analyse Ultrametrique, 9e Annee: 1981/82, No.1, Expose No.10, 12 p. (1983)., 1983.